

**OLD DOMINION UNIVERSITY**

Department of Biological Sciences  
Old Dominion University, Norfolk, Virginia 23529

**BENTHIC BIOLOGICAL MONITORING PROGRAM OF THE ELIZABETH RIVER  
WATERSHED (2004)**

Prepared by

Principal Investigator:

Dr. Daniel M. Dauer

Submitted to:

Mr. Roger Everton  
Virginia Department of Environmental Quality  
Tidewater Regional Office  
5636 Southern Blvd  
Virginia Beach, VA 23462

August, 2005

## Table of Contents

|   |      |
|---|------|
| EXECUTIVE SUMMARY .....   | 1    |
| INTRODUCTION .....  | 3    |
| RATIONALE .....   | 3    |
| METHODS .....   | 4    |
| Strata Sampled .....  | 4    |
| Probability-Based Sampling .....  | 4    |
| Probability-Based Estimation of Degradation .....   | 4    |
| Fixed-Point Station sampling .....  | 5    |
| Laboratory Analysis .....   | 5    |
| Benthic Index of Biotic Integrity .....   | 6    |
| B-IBI and Benthic Community Status Designations .....   | 6    |
| Further Information concerning the B-IBI .....  | 6    |
| Fixed-Point Stations in the Elizabeth River from the Chesapeake Bay Program .....   | 7    |
| Water Quality of the Elizabeth River .....  | 8    |
| RESULTS .....   | 8    |
| Benthic Community Condition using Probability-Based Sampling .....  | 8    |
| Environmental Parameters .....  | 8    |
| Benthic Community Condition .....   | 8    |
| Benthic Community Trends using Fixed -Point Stations .....  | 9    |
| Environmental Parameters .....  | 9    |
| Benthic Community .....   | 9    |
| Summary Patterns in the B-IBI and Selected Metrics .....  | 9    |
| Mainstem .....  | 9    |
| Southern Branch .....   | 10   |
| Western and Eastern Branches .....  | 11   |
| Lafayette River .....   | 11   |
| Benthic Community Trends in the James River of the Chesapeake Bay Program .....   | 12   |
| Water Quality of the Elizabeth River .....  | 12   |
| Discussion .....  | 13   |
| Watershed Level Condition of Benthic Communities .....  | 13   |
| Long-term Trends of Benthic Communities .....   | 13   |
| Water Quality of the Elizabeth River .....  | 14   |
| REFERENCES .....  | 15   |
| Glossary of selected terms .....  | 19   |
| Figures .....   | 20   |
| Tables .....  | 31   |
| Appendix A: Metrics and thresholds for calculating the Benthic Index Biotic Integrity .....   | A-41 |
| Appendix B: Summary of water quality status and trends for the Elizabeth River .....  | B-45 |
| Appendix C: Summary of benthic community status and trends for the Elizabeth River<br>stations (SBE2 and SBE5) of the Benthic Monitoring Program of the Chesapeake Bay<br>Program ..... | C-51 |
| Appendix D: Plots of fixed -point station B-IBI and metric values from 1999-2003 for the 14 stations<br>of the Elizabeth River Benthic Monitoring Program. ....                         | D-63 |

## EXECUTIVE SUMMARY

Macroenthic communities of the Elizabeth River watershed have been quantitatively sampled since summer 1999. This report presents the data from the sixth year of sampling in 2004. The three objectives of the Benthic Biological Monitoring Program of the Elizabeth River watershed are: (1) To characterize the health of the tidal waters of the Elizabeth River watershed as indicated by the structure of the benthic communities. (2) To conduct trend analyses on long-term data at 14 fixed-point stations to relate temporal trends in the benthic communities to changes in water and/or sediment quality. Trend analyses will be updated annually as new data are available. (3) To produce an historical data base that will allow annual evaluations of biotic impacts by comparing trends in status within probability-based strata and trends at fixed-point stations to changes in water and/or sediment quality.

The health of the benthic communities of the Elizabeth River watershed is characterized by combining the Benthic Index of Biotic Integrity (B-IBI) developed for the Chesapeake Bay and probability-based sampling. A probability-based sampling design allows calculation of confidence intervals around estimates of condition of the benthic communities and allows estimates of the areal extent of degradation of the benthic communities. The 2004 areal estimate of degraded benthic bottom was the highest level recorded since the beginning of this program in 1999. Based upon probability-based sampling the estimate of benthic bottom not meeting the benthic restoration goals was  $64 \pm 10.1\%$  in 1999,  $72 \pm 17.6\%$  in 2000,  $52 \pm 19.6\%$  in 2001,  $72 \pm 17.6\%$  in 2002,  $80 \pm 15.7\%$  in 2003 and  $88 \pm 12.7\%$  in 2004. Average B-IBI values for the Elizabeth River watershed were 2.7, 2.6, 2.7, 2.4, 2.3 and 2.2 respectively for the years 1999-2004.

Trend analyses were conducted using the data from the 14 fixed point stations for the period 1999-2004. Four stations showed trends in the B-IBI at  $p < 0.05$  with three stations showing improvements (LFA1, LFB1, ELD1) and one station deterioration (EBB1). Using the approach of the Chesapeake Bay Program, the status of each of the 14 fixed-point stations was characterized using the median value of the B-IBI for the last three years (2002-2004). Only one station, ELD1 in the Mainstem, had a B-IBI value over 3.0 and was considered to have met the Benthic Restoration Goals of the B-IBI. Two stations had a marginal status value of the B-IBI - a station in the Mainstem, ELD1, and the southernmost station in the Southern Branch, SBD4. All other fixed-point stations had a degraded or severely degraded category.

In general for the Elizabeth River watershed, species diversity and biomass remain below reference condition levels while abundance was often above reference condition levels and considered excessive. Community composition was unbalanced with levels of pollution indicative species above, and levels of pollution sensitive species, below reference conditions.

The water quality of the Elizabeth River can be generally characterized as follows: (1) nutrients have a poor status indicating high concentration levels, (2) there were improvements in long-term trends in surface total nitrogen levels (STN) and inorganic nitrogen levels (SDIN), and

(3) widespread improvements in long-term trends in surface total phosphorus levels (STP). Nutrient levels of the Elizabeth River exceed those of the lower section of the James River (Table 9). Nitrogen levels are highest in the Southern Branch with smaller differences between the branches of the river for phosphorus levels. The nutrient level in the Elizabeth River are more comparable to levels in the upper reaches of the James River in oligohaline and tidal freshwater regions (Dauer et al. 2003a,b; 2005). Chlorophyll levels, indicative of algal blooms when high, are good in both the Eastern Branch and Southern Branch in spite of high nutrient levels and good water clarity (Appendix B, Fig. B3). Chlorophyll levels are fair in the Western Branch but there is an improving long-term trend. Bottom dissolved oxygen are fair to good in all branches.



## INTRODUCTION

A long-term monitoring program of the macrobenthic communities of the Elizabeth River watershed was initiated in summer 1999. The three objectives of the Benthic Biological Monitoring Program of the Elizabeth River watershed are: (1) To characterize the health of the tidal waters of the Elizabeth River watershed as indicated by the structure of the benthic communities. This characterization is based upon application of benthic restoration goals and the Benthic Index of Biotic Integrity (B-IBI) developed for the Chesapeake Bay to the Elizabeth River Watershed (Ranasinghe et al. 1994; Weisberg et al. 1997; Alden et al. 2002). In each year 25 samples are randomly allocated in a probability-based sampling design. A probability-based sampling design allows calculation of confidence intervals around estimates of condition of the benthic communities. (2) To conduct trend analyses on long-term data at 14 fixed-point stations to relate temporal trends in the benthic communities to changes in water and/or sediment quality. Trend analyses will be updated annually as new data are available. (3) To produce an historical data base that will allow annual evaluations of biotic impacts by comparing trends in status within probability-based strata and trends at fixed-point stations to changes in water and/or sediment quality.

The macrobenthic communities of the Elizabeth River have been studied since the 1969 sampling of Boesch (1973) with three stations in the Mainstem of the river. Other important studies were limited to the Southern Branch of the river including seasonal sampling at 10 sites in 1977-1978 (Hawthorne and Dauer 1983), seasonal sampling at the same 10 sites a decade later in 1987-1988 by Hunley (1993), the establishment of two long-term monitoring stations in 1989 as part of the Virginia Chesapeake Bay Benthic Monitoring Program (Dauer et al. 1999) and summarizations of the two Southern Branch long-term monitoring stations (Dauer 1993; Dauer et al. 1993). The condition of the benthic community of the Elizabeth River watershed was characterized by spatially extensive sampling of the river in 1999 with 175 locations sampled among seven strata (Dauer 2000; Dauer and Llansó 2003). Beginning in 2000 the Elizabeth River watershed was sampled as a single stratum with the benthic community condition characterized at 25 random locations (Dauer 2001, 2002, 2003, 2004). This study updates the benthic community characterization of the Elizabeth River watershed base upon data collected in 2004.

## RATIONALE

Benthic invertebrates are used extensively as indicators of estuarine environmental status and trends because numerous studies have demonstrated that benthos respond predictably to many kinds of natural and anthropogenic stress (Pearson and Rosenberg 1978; Dauer 1993; Tapp et al. 1993; Wilson and Jeffrey 1994). Many characteristics of benthic assemblages make them useful indicators (Bilyard 1987), the most important of which are related to their exposure to stress and the diversity of their responses to stress. Exposure to hypoxia is typically greatest in near-bottom waters and anthropogenic contaminants often accumulate in sediments where benthos live. Benthic organisms generally have limited mobility and cannot avoid these adverse

conditions. This immobility is advantageous in environmental assessments because, unlike most pelagic fauna, benthic assemblages reflect local environmental conditions (Gray 1979). The structure of benthic assemblages responds to many kinds of stress because these assemblages typically include organisms with a wide range of physiological tolerances, life history strategies, feeding modes, and trophic interactions (Pearson and Rosenberg 1978; Rhoads et al. 1978; Boesch and Rosenberg 1981; Dauer 1993). Benthic community condition in the Chesapeake Bay watershed has been related in a quantitative manner to water quality, sediment quality, nutrient loads, and land use patterns (Dauer et al. 2000).

## **METHODS**

A glossary of selected terms used in this report is found on page 18.

### ***Strata Sampled***

In the summer of 1999, the Elizabeth River watershed was divided into five primary strata - the Mainstem of the river, the Lafayette River, the Southern Branch, Western Branch and Eastern Branch (Fig. 1). In addition two small creeks of the Southern Branch of the river were also sampled as part of a sediment contaminant remediation effort - Scuffletown Creek and Jones-Gilligan Creek. Beginning in 2000 and in subsequent years the Elizabeth River was sampled as a single stratum of 25 random samples. In 2001 Paradise Creek was sampled as a separate stratum.

### ***Probability-based Sampling***

Sampling design and methodologies for probability-based sampling are based upon procedures developed by EPA's Environmental Monitoring and Assessment Program (EMAP, Weisberg et al. 1993) and allow unbiased comparisons of conditions between strata (Dauer and Llansó 2003).

Within each probability-based stratum, 25 random locations were sampled using a 0.04 m<sup>2</sup> Young grab. The minimum acceptable depth of penetration of the grab was 7 cm. At each station one grab sample was taken for macrobenthic community analysis and an additional grab sample for sediment particle size analysis and the determination of total volatile solids. A 50 g subsample of the surface sediment was taken for sediment analyses. Salinity, temperature and dissolved oxygen were measured at the bottom and water depth was recorded.

### ***Probability-Based Estimation of Degradation***

Areal estimates of degradation of benthic community condition within a stratum can be made because all locations in each stratum are randomly selected. The estimate of the proportion of a stratum failing the Benthic Restoration Goals developed for Chesapeake Bay (Ranasinghe et al. 1994; updated in Weisberg et al. 1997) is the proportion of the 25 samples with an B-IBI value of less than 3.0. The process produces a binomial distribution: the percentage of the

stratum attaining goals versus the percentage not attaining the goals. With a binomial distribution the 95% confidence interval for these percentages can be calculated as:

$$95\% \text{ Confidence Interval} = p \pm 1.96 (\text{SQRT}(pq/N))$$

where p = percentage attaining goal, q = percentage not attaining goal and N = number of samples. This interval reflects the precision of measuring the level of degradation and indicates that with a 95% certainty the true level of degradation is within this interval. Differences between levels of degradation using a binomial distribution can be tested using the procedure of Schenker and Gentleman (2001).

For each stratum, 50 random points were selected using the GIS system of Versar, Inc. Decimal degree reference coordinates were used with a precision of 0.000001 degrees (approximately 1 meter) which is a smaller distance than the accuracy of positioning; therefore, no area of a stratum is excluded from sampling and every point within a stratum has a chance of being sampled. In the field the first 25 acceptable sites are sampled. Sites may be rejected because of inaccessibility by boat, inadequate water depth or inability of the grab to obtain an adequate sample (e.g., on hard bottoms).

### ***Fixed-Point Station Sampling***

Fourteen fixed point stations were established for long-term trend analysis (Fig. 2). All field collection procedures were the same as for probability based sampling except that three replicate Young grab sample were collected for macrobenthic community analysis.

### ***Laboratory Analysis***

Each replicate was sieved on a 0.5 mm screen, relaxed in dilute isopropyl alcohol and preserved with a buffered formalin-rose bengal solution. In the laboratory each replicate was sorted and all the individuals identified to the lowest possible taxon and enumerated. Biomass was estimated for each taxon as ash-free dry weight (AFDW) by drying to constant weight at 60 °C and ashing at 550 °C for four hours. Biomass was expressed as the difference between the dry and ashed weight.

Particle-size analysis was conducted using the techniques of Folk (1974). Each sediment sample is first separated into a sand fraction (> 63 µm) and a silt-clay fraction (< 63 µm). The sand fraction was dry sieved and the silt-clay fraction quantified by pipette analysis. For random stations, only the percent sand and percent silt-clay fraction were estimated. For the fixed-point stations particle-size distribution parameters were determined by the graphic and moment measures methods of Folk (1974). Total volatile solids of the sediment was estimated by the loss upon ignition method as described above and presented as percentage of the weight of the sediment.

## *Benthic Index of Biotic Integrity*

### **B-IBI and Benthic Community Status Designations**

The B-IBI is a multiple-metric index developed to identify the degree to which a benthic community meets the Chesapeake Bay Program's Benthic Community Restoration Goals (Ranasinghe et al. 1994; Weisberg et al. 1997; Alden et al. 2002). The B-IBI provides a means for comparing relative condition of benthic invertebrate communities across habitat types. It also provides a validated mechanism for integrating several benthic community attributes indicative of community health into a single number that measures overall benthic community condition.

The B-IBI is scaled from 1 to 5, and sites with values of 3 or more are considered to meet the Restoration Goals. The index is calculated by scoring each of several attributes as either 5, 3, or 1 depending on whether the value of the attribute at a site approximates, deviates slightly from, or deviates strongly from the values found at reference sites in similar habitats, and then averaging these scores across attributes. The criteria for assigning these scores are numeric and dependent on habitat type. Application of the index is limited to a summer index period from July 15th through September 30th. Habitat specific metrics and scoring thresholds are presented in Appendix A.

Benthic community condition was classified into four levels based on the B-IBI. Values  $\geq 2$  were classified as **severely degraded**; values from 2.1 to 2.6 were classified as **degraded**; values greater than 2.6 but less than 3.0 were classified as **marginal**; and values of 3.0 or more were classified as **meeting the goal**. Values in the marginal category do not meet the Restoration Goals, but they differ from the goals within the range of measurement error typically recorded between replicate samples. These categories are used in annual characterizations of the condition of the benthos in the Chesapeake Bay (Ranasinghe et al. 1994; Dauer et al. 1998a, 1998b; Ranasinghe et al. 1998; Dauer et al. 2002a,b; Llansó et al 2004).

### **Further Information concerning the B-IBI**

The analytical approach used to develop the B-IBI was similar to the one Karr et al. (1986) used to develop comparable indices for freshwater fish communities. Selection of benthic community metrics and metric scoring thresholds were habitat-dependent but by using categorical scoring comparisons between habitat types were possible. A six-step procedure was used to develop the index: (1) acquiring and standardizing data sets from a number of monitoring programs, (2) temporally and spatially stratifying data sets to identify seasons and habitat types, (3) identifying reference conditions, (4) selecting benthic community metrics, (5) selecting metric thresholds for scoring, and (6) validating the index with an independent data set (Weisberg et al. 1997). The B-IBI developed for Chesapeake Bay is based upon subtidal, unvegetated, infaunal macrobenthic communities. Hard-bottom communities, e.g., oyster beds, were not sampled because the sampling gears could not obtain adequate samples to characterize the associated infaunal communities. Infaunal communities associated with submerged aquatic vegetation

(SAV) were not avoided, but were rarely sampled due to the limited spatial extent of SAV in Chesapeake Bay.

Only macrobenthic data sets based on processing with a sieve of 0.5 mm mesh aperture and identified to the lowest possible taxonomic level were used. A data set of over 2,000 samples collected from 1984 through 1994 was used to develop, calibrate and validate the index (see Table 1 in Weisberg et al. 1997). Because of inherent temporal sampling limitations in some of the data sets, only data from the period of July 15 through September 30 were used to develop the index. A multivariate cluster analysis of the biological data was performed to define habitat types. Salinity and sediment type were the two important factors defining habitat types and seven habitats were identified - tidal freshwater, oligohaline, low mesohaline, high mesohaline sand, high mesohaline mud, polyhaline sand and polyhaline mud habitats (see Table 5 in Weisberg et al. 1997 and Appendix A of this report).

Reference conditions were determined by selecting samples which met all three of the following criteria: no sediment contaminant exceeded Long et al.'s (1995) effects range-median (ER-M) concentration, total organic content of the sediment was less than 2%, and bottom dissolved oxygen concentration was consistently high.

A total of 11 metrics representing measures of species diversity, community abundance and biomass, species composition, depth distribution within the sediment, and trophic composition were used to create the index (see Appendix). The habitat-specific metrics were scored and combined into a single value of the B-IBI. Thresholds for the selected metrics were based on the distribution of values for the metric at the reference sites. Data used for validation were collected between 1992 and 1994 and were independent of data used to develop the index. The B-IBI classified 93% of the validation sites correctly (Weisberg et al. 1997).

In tables presenting B-IBI results salinity classes are coded as follows: 1- tidal freshwater, 2 - oligohaline, 3- low mesohaline, 4 - high mesohaline and 5 - polyhaline. The two sediment classes are as follows: 1 - silt clay content < 40% and 2 - silt clay content  $\geq$  40%. All abundance values are individuals per m<sup>2</sup>, biomass values are AFDW g per m<sup>2</sup>, and pollution indicative, pollution sensitive and carnivore/omnivore metrics are percent of abundance or biomass as indicated in tables.

### ***Fixed-Point Stations in the Elizabeth River from the Chesapeake Bay Program***

Data concerning benthic community status and trends for James River, including the Elizabeth River, are collected by the Virginia Benthic Monitoring Program as part of the Chesapeake Bay Restoration Program. These data have been updated recently to include all data through 2004 and are presented here to summarize patterns for the five James River stations and two Elizabeth River stations that are part of that program. Details of collection and laboratory methodology can be found in Dauer et al. 2005 which can be downloaded in pdf format from the Old Dominion University Chesapeake Bay Program website <[www.chesapeakebay.odu.edu](http://www.chesapeakebay.odu.edu)> under "Reports." The James River Report includes the Elizabeth River, the Chickahominy

River and the Appomattox River.

### ***Water Quality of the Elizabeth River***

Data concerning water quality status and trends for the Elizabeth River are collected by the Virginia Water Quality Monitoring Program as part of the Chesapeake Bay Restoration Program. These data have been updated recently to include all data through 2004 and are presented here to summarize water quality of the Elizabeth River. Details of collection and laboratory methodology can be found in Dauer et al. 2005 which can be downloaded in pdf format from the Old Dominion University Chesapeake Bay Program website <[www.chesapeakebay.odu.edu](http://www.chesapeakebay.odu.edu)> under “Reports.” The James River Report includes the Elizabeth River, the Chickahominy River and the Appomattox River. Also available at this website are appendices that include (1) tables of status for all parameters measured at all stations sampled by each program, (2) tables of all parameters and metrics for which there was a significant trend, and (3) scatter plots of all parameters over time. There are five appendices: water quality, phytoplankton, primary productivity, zooplankton and benthos.

## **RESULTS**

### ***Benthic Community Condition using Probability-Based Sampling***

#### **Environmental Parameters**

All physical, chemical and sedimentary parameters are summarized in Table 1. Water depths varied from less than 1m to 17m reflecting shoal and channel depths. In 1999-2002 most stations had salinity values in the polyhaline range. During the summer of 2003 (Dauer 2004) and again in the 2004 collections most salinity values were in the high mesohaline range. Two stations in the Southern Branch had bottom dissolved oxygen measurements below 2.0 ppm. Macrobenthic communities are generally not altered by low dissolved oxygen unless values fall below at least 2.0 ppm (Diaz and Rosenberg 1995). As in previous collection years silt-clay content varied widely from less than 5% to greater than 95% and total volatile solids values were also less than 3%.

#### **Benthic Community Condition**

Benthic community parameters including the B-IBI value, abundance, biomass, Shannon diversity and selected metrics are summarized by station in Table 2. The average B-IBI values for the 25 random sites was 2.2. The distribution of the random sites and benthic community condition designations are shown in Figure 3. For the 2004 data,  $88 \pm 12.7$  % of the watershed had degraded benthos. Individual metric scores incorporated in the B-IBI are presented in Table 3. The dominant taxa of the random sites are summarized in Table 4. The pollution sensitive bivalve *Macoma balthica* was no longer listed as a density dominant as in 2003.

The B-IBI value, Shannon's index, abundance, biomass and the proportion of pollution sensitive and pollution indicative species for 1999-2004 are shown in Figs. 4-9. The 1999 value is the area-weighted average for the five strata sampled in that year.

### ***Benthic Community Trends using Fixed -Point Stations***

#### **Environmental Parameters**

All physical, chemical and sedimentary parameters are summarized in Table 5.

#### **Benthic Community**

Benthic community parameters including the B-IBI value, abundance, biomass, Shannon diversity and selected metrics are summarized by station in Table 6. Figure 7 lists the status of the 14 fixed-point stations. Status is determined by the three year (2002-2004) average B-IBI values at each station. Only one station, ELD1 in the Mainstem, had a B-IBI value over 3.0 and was considered to have met the Benthic Restoration Goals of the B-IBI. Two stations had a marginal status value of the B-IBI - a station in the Mainstem, ELC1, and the southernmost station in the Southern Branch, SBD4. All other fixed-point stations had a degraded or severely degraded category.

Four stations showed a trend in the B-IBI at  $p < 0.05$  with three stations showing improvements (LFA1, LFB1, ELD1) and one station deterioration (EBB1). Of the 17 significant trends in individual B-IBI metrics, 5 were degrading trends and 12 were improving trends. Of the five degrading trends, two were in the diversity index in the Southern Branch (SBEC1, SBED4), two were changes in community composition (a decrease in Pollution Sensitive Biomass at ELC1 and increase in Pollution Indicative Biomass at LFA1) and an excessive increase in abundance in the Southern Branch Station SBA1.

#### **Summary Patterns in the B-IBI and Selected Metrics**

##### **Mainstem**

- ☐ **Benthic Index of Biotic Integrity:** From 1999-2004, the B-IBI values for stations ELC1 and ELD1 were generally close to or above 3.0. Station ELF1 was consistently below 3.0. There was a significantly improving trend at Station ELD1 due in part to reaching a peak value in 2004.
- ☐ **Species Diversity:** The mouth stations (ELC1 and ELD1) generally had higher diversity values with the index typically above 2.2, while Station ELF1 generally had lower values, generally below 2.2. Species diversity for the Mainstem stations were generally at their lowest values in the last two years (2003-2004). These species diversity patterns were consistent with the general pattern for the B-IBI.
- ☐ **Abundance:** For 1999 - 2002 the mouth stations (ELC1 and ELD1) generally had

community abundance values between 2,000 - 3,000 individuals  $\text{m}^{-2}$  with highest values in 2003 around 6,000 individuals  $\text{m}^{-2}$ . The data for 2004 indicate a decrease to previous values for the earlier years. Station ELF1 showed a more erratic pattern with higher values in 2000 and 2003 exceeding 5,000 individuals  $\text{m}^{-2}$  and a peak in 2004 exceeding 10,000 individuals  $\text{m}^{-2}$ . In general the Mainstem stations had the lowest abundances compared to the other branches of the river. For high salinity benthic habitat types, an abundance in excess of 5,000 - 8,000 individuals  $\text{m}^{-2}$  is considered excessive and results in a low score for this metric.

- ❑ **Biomass:** No patterns were obvious in biomass values. In all branches of the river most biomass values were around 1.0 g AFDW  $\text{m}^{-2}$  with some stations reaching their highest value in the last year or two. All values were generally considered to be at insufficient levels relative to the Benthic Restoration Goals.
- ❑ **Community Composition:** The mouth stations (ELC1 and ELD1) had a percent composition of Pollution Indicative Species Abundance that was generally less than 20% while Station ELF1 generally had values above 30%. As would be expected, the relative abundance of Pollution Sensitive Species was inversely related to the patterns for Pollution Indicative Species. This pattern is due to the large increase in the pollution indicative spionid polychaete *Streblospio benedicti* in recent years. When averaging these three stations as a group *S. benedicti* abundance was generally below 2,000 individuals  $\text{m}^{-2}$  from 1999 to 2002 and in 2003-2004 was over 10,000 individuals  $\text{m}^{-2}$ .

### Southern Branch

- ❑ **Benthic Index of Biotic Integrity:** The B-IBI values for all stations were consistently below 3.0 with the highest values at upstream station SBD4.
- ❑ **Species Diversity:** Compared to the Mainstem stations, species diversity values in the Southern Branch were more consistent with the upper Mainstem Station ELF1 and were generally lower than the two Mainstem Stations near the mouth of the Elizabeth River. Lowest species diversity values generally occurred in the three years (2002 - 2004).
- ❑ **Abundance:** Community abundance values were much higher than in the Mainstem stations and typically exceeded 10,000 individuals  $\text{m}^{-2}$ . Such an abundance is considered to be excessive relative to the Benthic Restoration Goals. Station SBA1 had highest values in the last three years (2002-2004); Station SBB1 had high values 2002 and 2003 followed by 2004 decline; the middle station SBC1 peaked in 2001 but retained high values in the last three years; and the two upper stations, SBD2 and SBD4, reached their highest values in 2003 with a decline in 2004.
- ❑ **Biomass:** No patterns were obvious in biomass values with most values less than 1.5 AFDW  $\text{g m}^{-2}$ . The farthest upstream stations generally had the lowest biomass. In all branches of the river most biomass values were around 1.0 g AFDW  $\text{m}^{-2}$  with some



stations reaching their highest value in the last year or two. All values were generally considered to be at insufficient levels relative to the Benthic Restoration Goals.

- ❑ **Community Composition:** After 1999 the level of Pollution Indicative Species Abundance declined and was mostly below 30%. There was a general increase in Pollution Sensitive Species after 1999 due primarily to increased abundances of the pollution sensitive polychaete *Mediomastus ambiseta*.

### Western and Eastern Branches

- ❑ **Benthic Index of Biotic Integrity:** The B-IBI values for the Western Branch were generally below 3.0 and slightly higher at the upper station WBB5. The Eastern Branch station was near 3.0 the in 1999-2000 and had lower values in recent years. The Eastern Branch station, EBB1, showed a significantly deteriorating trend.
- ❑ **Species Diversity:** Species diversity values were generally at lowest levels in the last three years (2002-2004).
- ❑ **Abundance:** Community abundance values were higher than in the Mainstem Stations and typically exceeded 7,000 to 10,000 individuals m<sup>-2</sup> particularly in the later years of 2001 through 2004. In these years each station was dominated by the pollution sensitive polychaete *Mediomastus ambiseta*.
- ❑ **Biomass:** Biomass values peaked at all stations in 2003 due to a high settlement of the bivalve *Macoma balthica*; however, all three stations declined in 2004 to pre-2003 levels. Consistent with other metrics the upper Western Branch station had higher values than the lower Western Branch station. In all branches of the river most biomass values were around 1.0 g AFDW m<sup>-2</sup> with some stations reaching their highest value in the last year or two. All values were generally considered to be at insufficient levels relative to the Benthic Restoration Goals.
- ❑ **Community Composition:** In general in both branches Pollution Indicative Species declined while Pollution Sensitive Species composition a pattern due to increased abundances of the pollution sensitive polychaete *Mediomastus ambiseta* and the pollution sensitive bivalve *Macoma balthica*.

### Lafayette River

- ❑ **Benthic Index of Biotic Integrity:** Both stations in the Lafayette River had significantly improving trends with Station LFA1 with slight increase while Station LFB1 showed a greater increase; however, both station B-IBI values remain below the Benthic Restoration Goals.
- ❑ **Species Diversity:** Species diversity values were higher at the lower station and declined

in the later years in the upper station.

- ❑ **Abundance:** Community abundance levels were more comparable to the Mainstem and these values were lower than the Southern Branch, Western Branch and Eastern Branch values. Abundance values were variable at the lower station and generally increased at the upper station. Values at both stations were in the range of 3,000 to 5,000 individuals  $m^{-2}$ , a range often resulting in the maximum B-IBI metric score of 5.
- ❑ **Biomass:** Biomass values increased at both stations to their highest levels in 2003 with a decline in 2004. The pattern was generally due to the 2003 settlement of the bivalve *Macoma balthica* and the lack of persistence of this species in 2004. In all branches of the river most biomass values were around 1.0 g AFDW  $m^{-2}$  with some stations reaching their highest value in the last year or two. All values were generally considered to be at insufficient levels relative to the Benthic Restoration Goals.
- ❑ **Community Composition:** Pollution Indicative Species Abundance increased slightly in the lower station through 2003 with a decline in 2004, while at the upper station this metric has continued to increase. In contrast, Pollution Sensitive Species Abundance pattern was erratic in the lower station with a peak value in 2004 and increased greatly in the upper station.

### ***Benthic Community Trends in the James River of the Chesapeake Bay Program***

There are two stations in the Southern Branch of the Elizabeth River that are sampled as part of the larger Chesapeake Bay Program (SBE2 and SBE5, see Appendix C, Fig. C1). The status of both stations is poor but there are many improving trends in the benthic community and the B-IBI values at Station SBE5 shows a significant overall improving trend (Appendix C, Fig. C2, C4).

### ***Water Quality of the Elizabeth River***

Nutrient levels in all branches of the Elizabeth River are characterized by the Chesapeake Bay Program criteria as having a fair to poor status; however, there are several improving trends (Appendix B, Fig. B2 and Tables B1, B3). Surface and bottom nitrogen (STN and BTN) showed improving trends in almost all branches. The previously widespread improvements in dissolved inorganic nitrogen in all branches is now limited trends in surface DIN in the Southern Branch and bottom DIN in the Eastern Branch. Total phosphorus and dissolved inorganic phosphorus improving in all branches except the mainstem. Chlorophyll levels (SCHLA) had a good status in both the Southern Branch and the Eastern Branch, and fair status in the Mainstem and the Western Branch (Appendix B, Fig. B3). Indicators of water clarity, total suspended solids (STSS, BTSS) and secchi depth (SECCHI) generally showed improving trends in all branches. Finally, previously reported improving trends in bottom dissolved oxygen were no longer significant but dissolved oxygen status remains good to fair (Appendix B, Fig. B3, Table B3).

## Discussion

### *Watershed Level Condition of Benthic Communities*

Probability-based sampling allows an annual characterization of the overall condition of the benthic communities of the Elizabeth River watershed. In 1999 the condition of the macrobenthic communities of the Elizabeth River watershed was characterized for five strata consisting of the Mainstem of the River, the Lafayette River, the Southern Branch, Western Branch and Eastern Branch (Dauer 2000). The 1999 intensive sampling serves as a benchmark for all future analyses. The five strata were characterized in terms of benthic community condition into three categories: (1) the best condition in the Mainstem of the river, (2) the worst condition in the Southern Branch, and (3) intermediate condition in the Eastern Branch, Western Branch and Lafayette River. The Mainstem of the river had the highest average B-IBI value of 2.9, the Southern Branch the lowest value of 2.0 and the other branches had values between 2.5 and 2.7 with an overall average of 2.5. In 1999 each of the five strata were sampled at 25 random locations for a total of 125 random samples. In succeeding years the entire Elizabeth River watershed has been sampled as a single stratum of 25 random samples.

In 2004 the average watershed-level value for the B-IBI was the lowest recorded since 1999 and the area of benthic not meeting the Chesapeake Bay Benthic Restoration Goals was the highest recorded since 1999. Average B-IBI values for the Elizabeth River watershed were 2.2 (2004), 2.3 (2003), 2.4 (2002), 2.7 (2001), 2.6 (2000), 2.7 (1999) (see Dauer and Rodi 1999; Dauer 2000, 2001, 2002, 2003). Based upon probability-based sampling the estimate of benthic bottom not meeting the benthic restoration goals was  $88 \pm 12.7\%$  in 2004,  $80 \pm 15.7\%$  in 2003,  $76 \pm 16.7\%$  in 2002,  $52 \pm 19.6\%$  in 2001,  $72 \pm 17.6\%$  in 2000, and  $64 \pm 10.1\%$  in 1999.

Compared to the Chesapeake Bay Benthic Restoration Goals the macrobenthic communities of the Elizabeth River can be characterized as (1) having lower than expected species diversity and biomass, (2) abundance levels generally higher than reference conditions and (3) species composition with levels of pollution indicative species higher than reference conditions and levels of pollution sensitive species lower than reference conditions (Table 2; Figs. 4-9). However, there are some positive indications with the increase in the proportion of pollution sensitive species (Fig. 9).

### *Long-term trends of Benthic Communities*

Long-trend analyses in values of the B-IBI were conducted for the period 1999-2004. Four stations showed a trend in the B-IBI at  $p < 0.05$  with three stations showing improvements (LFA1, LFB1, ELD1) and one station deterioration (EBB1). Of the 17 significant trends in individual B-IBI metrics, 5 were degrading trends and 12 were improving trends. Of the five degrading trends, two were in the diversity index in the Southern Branch (SBEC1, SBED4), two were changes in community composition (a decrease in Pollution Sensitive Biomass at ELC1 and increase in Pollution Indicative Biomass at LFA1) and an excessive increase in abundance in the Southern Branch Station SBA1.ch (EBB1).

### ***Water Quality of the Elizabeth River***

The water quality of the Elizabeth River can be generally characterized as follows: (1) nutrients have a poor status indicating high concentration levels, (2) there were improvements in long-term trends in surface total nitrogen levels (STN) and inorganic nitrogen levels (SDIN), and (3) widespread improvements in long-term trends in surface total phosphorus levels (STP). Nutrient levels of the Elizabeth River exceed those of the lower section of the James River (Table 9). Nitrogen levels are highest in the Southern Branch with smaller differences between the branches of the river for phosphorus levels. The nutrient level in the Elizabeth River are more comparable to levels in the upper reaches of the James River in oligohaline and tidal freshwater regions (Dauer et al. 2003a,b; 2005). Chlorophyll levels, indicative of algal blooms when high, are good in both the Eastern Branch and Southern Branch in spite of high nutrient levels and good water clarity (Appendix B, Fig. B3). Chlorophyll levels are fair in the Western Branch but there is an improving long-term trend. The status of bottom dissolved oxygen was fair to good in all branches.

## REFERENCES

- Alden, R.W. III, D.M. Dauer, J.A. Ranasinghe, L.C. Scott, and R.J. Llansó. 2002. Statistical Verification of the Chesapeake Bay Benthic Index of Biotic Integrity. *Environmetrics* 13: 473- 498.
- Bilyard, G. R. 1987. The value of benthic infauna in marine pollution monitoring studies. *Marine Pollution Bulletin* 18:581-585.
- Boesch, D.F. 1973. Classification and community structure of macrobenthos in the Hampton Roads area, Virginia. *Marine Biology* 21: 226-244.
- Boesch, D. F. and R. Rosenberg. 1981. Response to stress in marine benthic communities, p. 179-200. In G. W. Barret and R. Rosenberg (eds.), Stress Effects on Natural Ecosystems. John Wiley & Sons, New York.
- Dauer, D.M. 1993. Biological criteria, environmental health and estuarine macrobenthic community structure. *Marine Pollution Bulletin* 26: 249-257.
- Dauer, D.M. 2000. Benthic Biological Monitoring Program of the Elizabeth River Watershed (1999). Final Report to the Virginia Department of Environmental Quality, Chesapeake Bay Program, 73 pp.
- Dauer, D.M. 2001. Benthic Biological Monitoring Program of the Elizabeth River Watershed (2000). Final Report to the Virginia Department of Environmental Quality, Chesapeake Bay Program, 35 pp. Plus Appendix.
- Dauer, D.M. 2002. Benthic Biological Monitoring Program of the Elizabeth River Watershed (2001) with a study of Paradise Creek. Final Report to the Virginia Department of Environmental Quality, Chesapeake Bay Program, 45 pp.
- Dauer, D.M. 2003. Benthic Biological Monitoring Program of the Elizabeth River Watershed (2002). Final Report to the Virginia Department of Environmental Quality, Chesapeake Bay Program, 56 pp.
- Dauer, D.M. 2004. Benthic Biological Monitoring Program of the Elizabeth River Watershed (2003). Final Report to the Virginia Department of Environmental Quality, Chesapeake Bay Program, 88 pp.
- Dauer, D.M., M. F. Lane, H.G. Marshall, K.E. Carpenter. 1998a. Status and trends in water quality and living resources in the Virginia Chesapeake Bay: 1985-1997. Final report to the Virginia Department of Environmental Quality. 84 pages.
- Dauer, D.M., M. F. Lane, H.G. Marshall, K.E. Carpenter and R.J. Diaz. 1999. Status and trends

- in water quality and living resources in the Virginia Chesapeake Bay: 1985-1998. Final report to the Virginia Department of Environmental Quality. 65 pages.
- Dauer, D.M. and R. J. Llansó. 2003. Spatial scales and probability based sampling in determining levels of benthic community degradation in the Chesapeake Bay. *Environmental Monitoring and Assessment* 81: 175-186.
- Dauer, D.M., M.W. Luchenback, and A.J. Rodi, Jr. 1993. Abundance biomass comparisons (ABC method): Effects of an estuary gradient, anoxic/hypoxic events and contaminated sediments. *Marine Biology* 116:507-518.
- Dauer, D.M., H.G. Marshall, K.E. Carpenter, J.R. Donat and M. F. Lane. 2003a. Status and trends in water quality and living resources in the Virginia Chesapeake Bay: James River (1985-2001). Final report to the Virginia Department of Environmental Quality. 108 pp.
- Dauer, D.M., H.G. Marshall, J.R. Donat, M. F. Lane, S.C. Doughten and F.A. Hoffman. 2003b. Status and trends in water quality and living resources in the Virginia Chesapeake Bay: James River (1985-2002). Final report to the Virginia Department of Environmental Quality. 94 pp.
- Dauer, D.M., H.G. Marshall, J.R. Donat, M. F. Lane, S.C. Doughten and F.A. Hoffman. 2005. Status and trends in water quality and living resources in the Virginia Chesapeake Bay: James River (1985-2003). Final report to the Virginia Department of Environmental Quality. 74 pp.
- Dauer, D.M., H.G. Marshall, K.E. Carpenter, M. F. Lane, R.W. Alden, III, K.K. Nesius and L.W. Haas. 1998b. Virginia Chesapeake Bay water quality and living resources monitoring programs: Executive Report, 1985-1996. Final report to the Virginia Department of Environmental Quality. 28 pages.
- Dauer, D.M., J. A. Ranasinghe, and S. B. Weisberg. 2000. Relationships between benthic community condition, water quality, sediment quality, nutrient loads, and land use patterns in Chesapeake Bay. *Estuaries* 23: 80-96.
- Dauer, D.M. and A.J. Rodi, Jr. 1999. Baywide benthic community condition based upon 1998 random probability based sampling. Final report to the Virginia Department of Environmental Quality. 126 pp.
- Diaz, R. J. and R. Rosenberg. 1995. Marine benthic hypoxia: a review of its ecological effects and the behavioural responses of benthic macrofauna. *Oceanography and Marine Biology Annual Review* 33:245-303.
- Folk, R.L. 1974. Petrology of sedimentary rocks. Hemphills, Austin, 170 pp.

- Gray, J. S. 1979. Pollution-induced changes in populations. *Transactions of the Royal Philosophical Society of London (B)* 286:545-561.
- Hawthorne, S.D. and D.M. Dauer. 1983. Macrobenthic communities of the lower Chesapeake Bay. III. Southern Branch of the Elizabeth River. *Internationale Revue der gesamten Hydrobiologie* 68: 193-205.
- Hunley, W.S. 1993. Evaluation of long term changes in the macrobenthic community of the Southern Branch of the Elizabeth River, Virginia. Master's Thesis. Old Dominion University. 120 pp.
- Karr, J. R., K. D. Fausch, P. L. Angermeier, P. R. Yant, and I. J. Schlosser. 1986. Assessing biological integrity in running waters: A method and its rationale. Special Publication 5. Illinois Natural History Survey, Champaign, Illinois.
- Llansó, R.J., F.S. Kelley and L.S. Scott. 2004. Long-term benthic monitoring and assessment component. Level I Comprehensive Report. July 1984 – December 2003. Final report to the Maryland Department of Natural Resources.
- Long, E. R., D. D. McDonald, S. L. Smith, and F. D. Calder. 1995. Incidence of adverse biological effects within ranges of chemical concentrations in marine and estuarine sediments. *Environmental Management* 19:81-95.
- Pearson, T. H. and R. Rosenberg. 1978. Macrobenthic succession in relation to organic enrichment and pollution of the marine environment. *Oceanography and Marine Biology Annual Review* 16:229-311.
- Ranasinghe, J.A., L.C. Scott and F.S. Kelley. 1998. Chesapeake Bay water quality monitoring program: Long-term benthic monitoring and assessment component, Level 1 Comprehensive Report (July 1984-December 1997). Prepared for the Maryland Department of Natural Resources by Versar, Inc., Columbia, MD.
- Ranasinghe, J.A., S.B. Weisberg, D.M. Dauer, L.C. Schaffner, R.J. Diaz and J.B. Frithsen. 1994. Chesapeake Bay benthic community restoration goals. Report for the U.S. Environmental Protection Agency, Chesapeake Bay Office and the Maryland Department of Natural Resources. 49 pp.
- Rhoads, D. C., P. L. McCall, and J. Y. Yingst. 1978. Disturbance and production on the estuarine sea floor. *American Scientist* 66:577-586.
- Schenker, N. and J.F. Gentleman. 2001. On judging the significance of differences by examining the overlap between confidence intervals. *The American Statistician* 55: 182-186
- Tapp, J. F., N. Shillabeer, and C. M. Ashman. 1993. Continued observation of the benthic fauna

of the industrialized Tees estuary, 1979-1990. *Journal of Experimental Marine Biology and Ecology* 172:67-80.

Weisberg, S.B., J.A. Ranasinghe, D.M. Dauer, L.C. Schaffner, R.J. Diaz and J.B. Frithsen. 1997. An estuarine benthic index of biotic integrity (B-IBI) for Chesapeake Bay. *Estuaries* 20: 149-158.

Wilson, J. G. and D. W. Jeffrey. 1994. Benthic biological pollution indices in estuaries, p. 311-327. In J. M. Kramer (ed.), *Biomonitoring of Coastal Waters and Estuaries*. CRC Press, Boca Raton, Florida.



## Glossary of selected terms

**Benthos** - refers to organisms that dwell on or within the bottom. Includes both hard substratum habitats (e.g. oyster reefs) and sedimentary habitats (sand and mud bottoms).

**B-IBI** - the benthic index of biotic integrity of Weisberg et al. (1997). The is a multi-metric index that compares the condition of a benthic community to reference conditions.

**Fixed Point Stations** - stations for long-term trend analysis whose location is unchanged over time.

**Habitat** - a local environment that has a benthic community distinct for other such habitat types. For the B-IBI of Chesapeake Bay seven habitat types were defined as combinations of salinity and sedimentary types - tidal freshwater, oligohaline, low mesohaline, high mesohaline sand, high mesohaline mud, polyhaline sand and polyhaline mud.

**Macrobenthos** - a size category of benthic organisms that are retained on a mesh of 0.5 mm.

**Metric** - a parameter or measurement of benthic community structure (e.g., abundance, biomass, species diversity).

**Probability based sampling** - all locations within a stratum have an equal chance of being sampled. Allows estimation of the percent of the stratum meeting or failing the benthic restoration goals.

**Random Station** - a station selected randomly within a stratum. In every succeeding sampling event new random locations are selected.

**Reference condition** - the structure of benthic communities at reference sites.


**Reference sites** - sites determined to be minimally impacted by anthropogenic stress. Conditions at theses sites are considered to represent goals for restoration of impacted benthic communities. Reference sites were selected by Weisberg et al. (1997) as those outside highly developed watersheds, distant from any point-source discharge, with no sediment contaminant effect, with no low dissolved oxygen effect and with a low level of organic matter in the sediment.

**Restoration Goal** - refers to obtaining an average B-IBI value of 3.0 for a benthic community indicating that values for metrics approximate the reference condition.


**Stratum** - a geographic region of unique ecological condition or managerial interest. In the 1999 study the primary strata were the Mainstem of the river, the Lafayette River, the Eastern Branch, Western Branch and Southern Branch. In succeeding years the entire Elizabeth River watershed was sampled as a single stratum.

**Threshold** - a value of a metric that determines the B-IBI scoring. For all metrics except abundance and biomass, two thresholds are used - the lower 5<sup>th</sup> percentile and the 50<sup>th</sup> percentile (median) of the distribution of values at reference sites. Samples with metric values less than the lower 5<sup>th</sup> percentile are scored as a 1. Samples with values between the 5<sup>th</sup> and 50<sup>th</sup> metrics are scored as 3 and values greater than the 50<sup>th</sup> percentile are scored as 5. For abundance and biomass, values below the 5<sup>th</sup> and above the 95<sup>th</sup> percentile are scored as 1, values between the 5<sup>th</sup> and 25<sup>th</sup> and the 75<sup>th</sup> and 95<sup>th</sup> percentiles are scored as 3 and values between the 25<sup>th</sup> and 75<sup>th</sup> percentiles are scored as 5.


## Figures




**Figure 1.** Elizabeth River watershed showing the five major segments sampled in 1999. Insert shows Scuffletown Creek and the Jones-Gilligan Creek strata also sampled in 1999 and Paradise Creek stratum sampled in 2001.



**Figure 2.** Elizabeth River watershed showing the 14 fixed-point stations for long-term trend analyses.




**Figure 3.** Random samples collected in 2004. Shown is the condition of the benthic communities using the B-IBI value.




**Figure 4.** Benthic Index of Biotic Integrity. Shown are mean values for the entire Elizabeth River watershed from the probability-based program. Dashed line indicates restoration goal.

## Shannon Diversity Index

Dashed lines indicate range of goals




**Figure 5.** Shannon-Weiner Diversity Index. Shown are mean values for the entire Elizabeth River watershed from the probability-based program. Dashed lines indicates range of restoration goals.




**Figure 6.** Abundance per m<sup>2</sup>. Shown are mean values for the entire Elizabeth River watershed from the probability-based program. Dashed lines indicates range of restoration goals.







**Figure 7.** AFDW biomass per m<sup>2</sup>. Shown are mean values for the entire Elizabeth River watershed from the probability-based program. Dashed lines indicates range of restoration goals.



**Figure 8.** Percentage of Pollution Indicative Species Abundance. Shown are mean values for the entire Elizabeth River watershed from the probability-based program. Dashed lines indicates range of restoration goals.



**Figure 9.** Percentage of Pollution Sensitive Species Abundance. Shown are mean values for the entire Elizabeth River watershed from the probability-based program. Dashed lines indicates range of restoration goals.



**Figure 11.** Status of and long-term trends in the Benthic IBI for the Elizabeth River Project monitoring stations for the period of 1999 through 2004. All trends shown were significant at  $p \leq 0.05$ .

## Tables

Table1. Random Stations of the Elizabeth River sampled in 2004.  
Summary of physical-chemical parameters.

| STATION | Date collected | LATITUDE | LONGITUDE | Water depth (m) | Temperature (°C) | Salinity (ppt) | Dissolved oxygen (ppm) | Silt_clay content (%) | Volatile organics (%) |
|---------|----------------|----------|-----------|-----------------|------------------|----------------|------------------------|-----------------------|-----------------------|
| 11Z01   | 8/10/04        | 36.91887 | -76.34747 | 3.3             | 26.1             | 16.0           | 6.2                    | 90.0                  | 2.3                   |
| 11Z02   | 8/10/04        | 36.91250 | -76.31136 | 1.0             | 26.2             | 15.1           | 7.7                    | 88.9                  | 2.5                   |
| 11Z03   | 8/10/04        | 36.90621 | -76.30534 | 6.0             | 26.0             | 15.0           | 6.4                    | 63.7                  | 2.4                   |
| 11Z04   | 8/10/04        | 36.90535 | -76.29982 | 3.3             | 25.9             | 14.4           | 7.0                    | 97.1                  | 2.7                   |
| 11Z05   | 8/10/04        | 36.90616 | -76.28863 | 1.5             | 27.7             | 13.9           | 9.3                    | 76.6                  | 1.3                   |
| 11Z07   | 8/10/04        | 36.88849 | -76.27766 | 2.0             | 25.8             | 12.6           | 4.6                    | 91.2                  | 2.8                   |
| 11Z08   | 8/10/04        | 36.89226 | -76.33105 | 4.8             | 26.5             | 15.9           | 5.6                    | 73.2                  | 1.7                   |
| 11Z10   | 8/10/04        | 36.87972 | -76.33909 | 9.3             | 25.3             | 20.8           | 3.6                    | 94.5                  | 2.5                   |
| 11Z11   | 8/10/04        | 36.87857 | -76.33682 | 11.4            | 25.2             | 21.2           | 3.6                    | 92.6                  | 2.1                   |
| 11Z12   | 8/10/04        | 36.87785 | -76.33335 | 17.0            | 25.0             | 23.0           | 3.0                    | 96.2                  | 2.3                   |
| 11Z13   | 8/10/04        | 36.87444 | -76.33789 | 8.0             | 25.4             | 20.6           | 3.4                    | 95.9                  | 2.4                   |
| 11Z15   | 8/17/04        | 36.84754 | -76.35846 | 4.0             | 25.3             | 15.4           | 3.7                    | 99.2                  | 2.5                   |
| 11Z16   | 8/17/04        | 36.83595 | -76.38295 | 1.5             | 25.2             | 9.9            | 3.4                    | 94.8                  | 2.5                   |
| 11Z17   | 8/10/04        | 36.85998 | -76.31700 | 15.4            | 25.1             | 22.3           | 3.0                    | 93.1                  | 2.4                   |
| 11Z18   | 8/18/04        | 36.78412 | -76.30659 | 6.0             | 26.3             | 15.4           | 1.8                    | 78.3                  | 3.3                   |
| 11Z19   | 8/18/04        | 36.75804 | -76.30816 | 1.0             | 24.9             | 1.2            | 4.5                    | 3.4                   | 0.6                   |
| 11Z20   | 8/18/04        | 36.75134 | -76.29599 | 10.0            | 26.8             | 17.4           | 0.7                    | 98.4                  | 3.6                   |
| 11Z21   | 8/18/04        | 36.73463 | -76.28695 | 5.5             | 27.0             | 10.0           | 2.5                    | 11.0                  | 1.0                   |
| 11Z23   | 8/18/04        | 36.84173 | -76.28990 | 12.3            | 25.1             | 18.9           | 2.0                    | 97.5                  | 2.6                   |
| 11Z26   | 8/10/04        | 36.89807 | -76.29249 | 2.4             | 25.8             | 13.8           | 6.8                    | 97.2                  | 2.3                   |
| 11Z27   | 8/17/04        | 36.85451 | -76.36059 | 1.0             | 27.3             | 10.8           | 5.8                    | 91.4                  | 2.0                   |
| 11Z28   | 8/10/04        | 36.86608 | -76.34327 | 0.8             | 27.7             | 15.8           | 11.4                   | 2.2                   | 0.2                   |
| 11Z30   | 8/18/04        | 36.82094 | -76.23964 | 1.0             | 27.2             | 1.5            | 6.7                    | 96.5                  | 3.3                   |
| 11Z31   | 8/18/04        | 36.83659 | -76.29430 | 14.5            | 25.0             | 18.8           | 2.0                    | 95.2                  | 2.4                   |
| 11Z32   | 8/18/04        | 36.83297 | -76.21110 | 2.0             | 27.2             | 4.2            | 6.2                    | 82.2                  | 2.9                   |

Table 2. Random Stations of the Elizabeth River sampled in 2004. Summary of benthic community parameters. Abundance reported in ind/m2, biomass reported as grams/m<sup>2</sup>, all other abundance and biomass metrics are percentages.

| Station | BIBI | Abundance | Biomass | Shannon Index | Pollution Indicative Abundance | Pollution Sensitive Abundance | Pollution Indicative Biomass | Pollution Sensitive Biomass | Carnivore Omnivore Abundance | Deep Deposit Feeder Abundance |
|---------|------|-----------|---------|---------------|--------------------------------|-------------------------------|------------------------------|-----------------------------|------------------------------|-------------------------------|
| 11Z01   | 3.00 | 3,697     | 0.726   | 2.90          | 14.1                           | 64.4                          | 31.3                         | 50.0                        | 31.3                         | 44.8                          |
| 11Z02   | 2.00 | 8,641     | 0.953   | 1.73          | 8.4                            | 70.9                          | 16.7                         | 31.0                        | 7.6                          | 84.0                          |
| 11Z03   | 1.67 | 7,076     | 0.567   | 1.76          | 23.1                           | 66.7                          | 48.0                         | 32.0                        | 8.3                          | 70.2                          |
| 11Z04   | 2.33 | 8,119     | 0.953   | 1.28          | 19.6                           | 76.0                          | 26.2                         | 64.3                        | 2.5                          | 77.4                          |
| 11Z05   | 1.67 | 7,348     | 0.476   | 1.21          | 11.1                           | 83.6                          | 9.5                          | 57.1                        | 4.9                          | 81.5                          |
| 11Z07   | 2.67 | 5,919     | 1.111   | 1.05          | 14.6                           | 82.0                          | 2.0                          | 83.7                        | 4.2                          | 79.7                          |
| 11Z08   | 3.33 | 2,019     | 0.295   | 2.36          | 34.8                           | 52.8                          | 23.1                         | 38.5                        | 57.3                         | 2.2                           |
| 11Z10   | 2.00 | 6,418     | 0.726   | 2.39          | 15.5                           | 69.3                          | 37.5                         | 37.5                        | 17.3                         | 64.7                          |
| 11Z11   | 1.67 | 3,810     | 0.794   | 2.31          | 19.6                           | 67.9                          | 51.4                         | 25.7                        | 20.2                         | 60.7                          |
| 11Z12   | 1.00 | 680       | 0.181   | 1.70          | 76.7                           | 13.3                          | 62.5                         | 25.0                        | 10.0                         | 20.0                          |
| 11Z13   | 2.00 | 6,532     | 0.748   | 2.39          | 16.3                           | 64.9                          | 48.5                         | 39.4                        | 20.1                         | 62.5                          |
| 11Z15   | 2.00 | 9,344     | 0.748   | 0.70          | 9.2                            | 89.1                          | 15.2                         | 54.5                        | 1.0                          | 89.3                          |
| 11Z16   | 2.20 | 13,086    | 0.748   | 0.64          | 6.4                            | 90.6                          | 24.2                         | 66.7                        | 0.9                          | 93.1                          |
| 11Z17   | 3.00 | 499       | 0.454   | 2.61          | 0.0                            | 63.6                          | 0.0                          | 80.0                        | 68.2                         | 4.5                           |
| 11Z18   | 2.33 | 3,039     | 0.318   | 1.94          | 3.7                            | 80.6                          | 21.4                         | 28.6                        | 32.8                         | 60.4                          |
| 11Z19   | 3.50 | 295       | 0.113   | 1.57          | 15.4                           | 53.8                          | 40.0                         | 40.0                        | 7.7                          | 0.0                           |
| 11Z20   | 2.67 | 17,305    | 4.604   | 1.11          | 12.3                           | 81.8                          | 13.3                         | 77.8                        | 6.9                          | 81.4                          |
| 11Z21   | 1.80 | 3,583     | 0.227   | 1.54          | 58.9                           | 39.2                          | 10.0                         | 70.0                        | 20.9                         | 20.3                          |
| 11Z23   | 1.67 | 8,165     | 1.179   | 1.72          | 18.3                           | 75.6                          | 44.2                         | 40.4                        | 7.8                          | 75.0                          |
| 11Z26   | 2.00 | 6,486     | 0.658   | 1.83          | 18.5                           | 56.6                          | 13.8                         | 34.5                        | 5.2                          | 75.9                          |
| 11Z27   | 1.80 | 8,959     | 1.225   | 1.19          | 13.9                           | 79.7                          | 1.9                          | 37.0                        | 4.8                          | 80.5                          |
| 11Z28   | 2.67 | 2,722     | 0.522   | 2.68          | 19.2                           | 39.2                          | 8.7                          | 21.7                        | 6.7                          | 45.0                          |
| 11Z30   | 1.67 | 34,927    | 1.565   | 1.58          | 41.8                           | 42.1                          | 20.3                         | 40.6                        | 0.6                          | 56.9                          |
| 11Z31   | 2.00 | 6,985     | 0.590   | 1.53          | 9.4                            | 77.9                          | 42.3                         | 34.6                        | 6.5                          | 84.7                          |
| 11Z32   | 1.00 | 13,744    | 1.361   | 1.17          | 18.8                           | 74.3                          | 1.7                          | 93.3                        | 1.2                          | 79.4                          |
| Mean    | 2.15 | 7,576     | 0.874   | 1.72          | 20.0                           | 66.2                          | 24.5                         | 48.2                        | 14.2                         | 59.8                          |
| SE      | 0.13 | 1,412     | 0.172   | 0.12          | 3.4                            | 3.7                           | 3.6                          | 4.1                         | 3.4                          | 5.7                           |

Table 3. Random Stations of the Elizabeth River sampled in 2004. Summary of benthic community parameters scores of the BIBI.

| Station | BIBI | Salinity Class | Sediment Class | Shannon Index | Abundance | Biomass | Pollution Indicative Abundance | Pollution Sensitive Abundance | Pollution Indicative Biomass | Pollution Sensitive Biomass | Carnivore Omnivore Abundance |
|---------|------|----------------|----------------|---------------|-----------|---------|--------------------------------|-------------------------------|------------------------------|-----------------------------|------------------------------|
| 11Z01   | 3.00 | 4              | 2              | 3             | 3         | 3       | 1                              | .                             | .                            | 3                           | 5                            |
| 11Z02   | 2.00 | 4              | 2              | 1             | 1         | 3       | 3                              | .                             | .                            | 3                           | 1                            |
| 11Z03   | 1.67 | 4              | 2              | 1             | 1         | 3       | 1                              | .                             | .                            | 3                           | 1                            |
| 11Z04   | 2.33 | 4              | 2              | 1             | 1         | 3       | 3                              | .                             | .                            | 5                           | 1                            |
| 11Z05   | 1.67 | 4              | 2              | 1             | 1         | 1       | 3                              | .                             | .                            | 3                           | 1                            |
| 11Z07   | 2.67 | 4              | 2              | 1             | 1         | 3       | 5                              | .                             | .                            | 5                           | 1                            |
| 11Z08   | 3.33 | 4              | 2              | 3             | 5         | 1       | 3                              | .                             | .                            | 3                           | 5                            |
| 11Z10   | 2.00 | 5              | 2              | 1             | 3         | 3       | 1                              | .                             | .                            | 3                           | 1                            |
| 11Z11   | 1.67 | 5              | 2              | 1             | 3         | 3       | 1                              | .                             | .                            | 1                           | 1                            |
| 11Z12   | 1.00 | 5              | 2              | 1             | 1         | 1       | 1                              | .                             | .                            | 1                           | 1                            |
| 11Z13   | 2.00 | 5              | 2              | 1             | 3         | 3       | 1                              | .                             | .                            | 3                           | 1                            |
| 11Z15   | 2.00 | 4              | 2              | 1             | 1         | 3       | 3                              | .                             | .                            | 3                           | 1                            |
| 11Z16   | 2.20 | 3              | .              | 1             | 1         | 1       | .                              | 5                             | .                            | 3                           | .                            |
| 11Z17   | 3.00 | 5              | 2              | 3             | 1         | 1       | 5                              | .                             | .                            | 3                           | 5                            |
| 11Z18   | 2.33 | 4              | 2              | 1             | 3         | 1       | 3                              | .                             | .                            | 1                           | 5                            |
| 11Z19   | 3.50 | 2              | .              | .             | 3         | .       | .                              | .                             | .                            | .                           | 1                            |
| 11Z20   | 2.67 | 4              | 2              | 1             | 1         | 5       | 3                              | .                             | .                            | 5                           | 1                            |
| 11Z21   | 1.80 | 3              | .              | 1             | 3         | 1       | .                              | 1                             | .                            | 3                           | .                            |
| 11Z23   | 1.67 | 5              | 2              | 1             | 1         | 3       | 1                              | .                             | .                            | 3                           | 1                            |
| 11Z26   | 2.00 | 4              | 2              | 1             | 1         | 3       | 3                              | .                             | .                            | 3                           | 1                            |
| 11Z27   | 1.80 | 3              | .              | 1             | 1         | 3       | .                              | 3                             | .                            | 1                           | .                            |
| 11Z28   | 2.67 | 4              | 1              | 3             | 5         | 1       | .                              | 3                             | 3                            | .                           | 1                            |
| 11Z30   | 1.67 | 2              | .              | .             | 1         | .       | .                              | .                             | .                            | .                           | 1                            |
| 11Z31   | 2.00 | 5              | 2              | 1             | 3         | 3       | 1                              | .                             | .                            | 3                           | 1                            |
| 11Z32   | 1.00 | 2              | .              | .             | 1         | .       | .                              | .                             | .                            | .                           | 1                            |



| Table 4. Random Stations of the Elizabeth River sampled in 2004.<br>Dominant tax by abundance. Taxon code: B = bivalve, C = cumacean, G = gastropod, I = isopod, N = nemertean, O = oligochaete, P = polychaete. |                                |                              |
|--|--------------------------------|------------------------------|
| Rank   | Taxon                          | Abundance per m <sup>2</sup> |
| 1  | Mediomastus ambiseta (P)       | 4,779                        |
| 2  | Streblospio benedicti (P)      | 1,261                        |
| 3  | Tubificoides spp. Group I (O)  | 477                          |
| 4  | Paraprionospio pinnata (P)     | 225                          |
| 5  | Glycinde solitaria (P)         | 191                          |
| 6  | Acteocina canaliculata (G)     | 142                          |
| 7  | Tubificoides heterochaetus (O) | 88                           |
| 8  | Podarkeopsis levifusca (P)     | 57                           |
| 9  | Cyathura polita (I)            | 47                           |
| 10   | Leitoscoloplos spp. (P)        | 36                           |
| 11   | Gemma gemma (B)                | 31                           |
| 12   | Nemertea spp. (N)              | 29                           |
| 13   | Edotea triloba (I)             | 27                           |
| 14   | Leucon americanus (C)          | 25                           |
| 15   | Neanthes succinea (P)          | 25                           |

| Table 5. Fixed Stations of the Elizabeth River sampled in 2004. Summary of physical-chemical parameters. |                |          |           |                    |                     |                   |                              |                          |                             |
|--|----------------|----------|-----------|--------------------|---------------------|-------------------|------------------------------|--------------------------|-----------------------------|
| STATION  | Date collected | LATITUDE | LONGITUDE | Water depth<br>(m) | Temperature<br>(°C) | Salinity<br>(ppt) | Dissolved<br>oxygen<br>(ppm) | Silt_clay<br>content (%) | Volatile<br>organics<br>(%) |
| EBB1   | 8/18/04        | 36.8378  | -76.2422  | 1.5                | 27.4                | 8.8               | 10.6                         | 69.0                     | 5.7                         |
| ELC1   | 8/17/04        | 36.8796  | -76.3476  | 3.1                | 25.2                | 16.1              | 4.1                          | 39.5                     | 1.8                         |
| ELD1   | 8/17/04        | 36.8614  | -76.3357  | 2.2                | 25.3                | 15.7              | 5.3                          | 3.0                      | 0.4                         |
| ELF1   | 8/18/04        | 36.8486  | -76.2967  | 11.0               | 25.3                | 18.5              | 2.2                          | 92.3                     | 7.6                         |
| LFA1   | 8/17/04        | 36.9092  | -76.3138  | 3.0                | 24.7                | 16.4              | 4.2                          | 77.8                     | 3.4                         |
| LFB1   | 8/17/04        | 36.8896  | -76.2830  | 3.5                | 25.0                | 14.9              | 3.3                          | 99.0                     | 6.7                         |
| SBA1   | 8/18/04        | 36.8255  | -76.2907  | 14.0               | 25.9                | 19.0              | 1.5                          | 96.6                     | 8.4                         |
| SBB1   | 8/18/04        | 36.8117  | -76.2886  | 2.0                | 27.7                | 11.6              | 3.5                          | 50.1                     | 6.7                         |
| SBC1   | 8/18/04        | 36.7994  | -76.2944  | 11.5               | 25.8                | 18.3              | 1.5                          | 87.1                     | 8.0                         |
| SBD1   | 8/18/04        | 36.7796  | -76.3106  | 10.2               | 26.2                | 18.0              | 1.2                          | 70.7                     | 7.9                         |
| SBD2   | 8/18/04        | 36.7668  | -76.2969  | 2.5                | 27.2                | 9.0               | 2.8                          | 77.4                     | 10.7                        |
| SBD4   | 8/18/04        | 36.7402  | -76.2990  | 2.0                | 27.6                | 9.9               | 3.0                          | 8.0                      | 0.9                         |
| WBB1   | 8/17/04        | 36.8462  | -76.3576  | 2.7                | 25.5                | 15.0              | 4.8                          | 91.4                     | 5.3                         |
| WBB5   | 8/17/04        | 36.8293  | -76.3932  | 1.5                | 24.7                | 7.8               | 3.6                          | 82.9                     | 5.6                         |

| Table 6. Fixed Point Stations of the Elizabeth River sampled in 2004. Summary of benthic community parameters. All values are station means (n=3). Abundance reported as ind/m <sup>2</sup> , biomass reported as grams/m <sup>2</sup> , all other abundance and biomass metrics are percentages. |      |           |         |               |                                |                               |                              |                             |                              |                               |
|---|------|-----------|---------|---------------|--------------------------------|-------------------------------|------------------------------|-----------------------------|------------------------------|-------------------------------|
| Station   | BIBI | Abundance | Biomass | Shannon Index | Pollution Indicative Abundance | Pollution Sensitive Abundance | Pollution Indicative Biomass | Pollution Sensitive Biomass | Carnivore Omnivore Abundance | Deep Deposit Feeder Abundance |
| EBB1  | 1.80 | 8996      | 0.990   | 1.28          | 28.7                           | 70.3                          | 30.8                         | 61.9                        | 3.3                          | 70.5                          |
| ELC1  | 3.00 | 2911      | 0.363   | 2.29          | 10.7                           | 71.3                          | 25.6                         | 36.3                        | 28.5                         | 59.7                          |
| ELD1  | 3.89 | 1920      | 0.355   | 3.06          | 7.1                            | 63.6                          | 16.7                         | 35.3                        | 34.7                         | 39.1                          |
| ELF1  | 1.67 | 11378     | 1.051   | 1.67          | 17.5                           | 71.4                          | 35.4                         | 45.8                        | 4.4                          | 78.5                          |
| LFA1  | 2.22 | 3644      | 0.733   | 2.15          | 14.0                           | 72.3                          | 38.0                         | 27.1                        | 14.2                         | 69.0                          |
| LFB1  | 2.44 | 10713     | 1.504   | 1.19          | 18.1                           | 76.0                          | 5.9                          | 81.6                        | 2.0                          | 79.0                          |
| SBA1  | 1.67 | 16314     | 1.058   | 1.53          | 12.1                           | 75.1                          | 34.4                         | 54.0                        | 4.5                          | 83.9                          |
| SBB1  | 2.33 | 1746      | 0.673   | 2.08          | 26.8                           | 63.4                          | 12.6                         | 42.8                        | 34.8                         | 62.2                          |
| SBC1  | 1.78 | 16314     | 0.990   | 1.07          | 10.2                           | 85.7                          | 34.8                         | 54.3                        | 2.6                          | 87.3                          |
| SBD1  | 1.56 | 2548      | 0.287   | 1.33          | 8.9                            | 84.5                          | 36.4                         | 29.4                        | 9.4                          | 80.1                          |
| SBD2  | 2.73 | 1822      | 0.219   | 1.89          | 20.2                           | 76.7                          | 13.7                         | 70.2                        | 48.2                         | 30.9                          |
| SBD4  | 2.87 | 2835      | 0.265   | 1.53          | 8.1                            | 88.2                          | 8.6                          | 59.9                        | 29.2                         | 60.0                          |
| WBB1  | 1.78 | 7053      | 0.476   | 1.03          | 15.3                           | 82.6                          | 25.5                         | 57.4                        | 3.8                          | 80.9                          |
| WBB5  | 1.67 | 9843      | 0.816   | 1.26          | 17.3                           | 75.9                          | 8.2                          | 55.6                        | 3.4                          | 78.8                          |

Table 7. Status in benthic community condition based on the Benthic IBI at the Elizabeth River Project monitoring stations for the period of 2001 through 2004.

| Station         | Mean IBI | Status            |
|-----------------|----------|-------------------|
| Mainstem        |          |                   |
| ELC1            | 2.9      | Marginal          |
| ELD1            | 3.0      | Meets Goals       |
| ELF1            | 1.9      | Severely degraded |
| Southern Branch |          |                   |
| SBA1            | 1.5      | Severely degraded |
| SBB1            | 2.3      | Degraded          |
| SBC1            | 2.0      | Severely degraded |
| SBD1            | 1.8      | Severely degraded |
| SBD2            | 2.3      | Degraded          |
| SBD4            | 2.7      | Marginal          |
| Western Branch  |          |                   |
| WBB1            | 2.1      | Degraded          |
| WBB5            | 2.3      | Degraded          |
| Eastern Branch  |          |                   |
| EBB1            | 2.1      | Degraded          |
| Lafayette River |          |                   |
| LFA1            | 2.2      | Degraded          |
| LFB1            | 2.5      | Degraded          |

Table 8. Significant long term trends in the B-IBI and associated metrics for the Elizabeth River Project monitoring stations for the period of 1999 through 2004. All trends shown were significant at  $p \leq 0.05$ .

| Station | Water Body               | Variable                               | P value | Baseline Mean | Slope   | % Change |
|---------|--------------------------|--|---------|---------------|---------|----------|
| ELC1    | Elizabeth River Mainstem | Pollution Sensitive Species Biomass    | 0.0500  | 47.51         | -3.79   | -19.0    |
| ELD1    | Elizabeth River Mainstem | Benthic Index of Biotic Integrity      | 0.0500  | 3.22          | 0.19    | 29.4     |
| ELF1    | Elizabeth River Mainstem | Pollution Sensitive Species Abundance  | 0.0500  | 24.27         | 11.32   | 56.6     |
| EBB1    | Eastern Branch           | Benthic Index of Biotic Integrity      | 0.0500  | 3.03          | -0.32   | -53.0    |
| EBB1    | Eastern Branch           | Pollution Sensitive Species Abundance  | 0.0500  | 16.07         | 14.95   | 74.8     |
| FA1     | Lafayette River          | Benthic Index of Biotic Integrity      | 0.0367  | 1.93          | 0.08    | 20.3     |
| FA1     | Lafayette River          | Pollution Indicative Species Biomass   | 0.0367  | 41.83         | 2.91    | 14.6     |
| FB1     | Lafayette River          | Benthic Index of Biotic Integrity      | 0.0267  | 1.97          | 0.18    | 45.8     |
| FB1     | Lafayette River          | Total Biomass per square meter         | 0.0267  | 0.49          | 0.26    | 269.8    |
| FB1     | Lafayette River          | Pollution Sensitive Species Abundance  | 0.0267  | 6.36          | 20.50   | 102.5    |
| FB1     | Lafayette River          | Pollution Sensitive Species Biomass    | 0.0267  | 22.21         | 18.76   | 93.8     |
| SBA1    | Southern Branch          | Total Abundance per square meter       | 0.0500  | 3863.16       | 3315.82 | 429.2    |
| SBB1    | Southern Branch          | Pollution Sensitive Species Abundance  | 0.0500  | 31.75         | 13.31   | 66.6     |
| SBC1    | Southern Branch          | Shannon-Weiner Diversity Index         | 0.0500  | 2.46          | -0.39   | -78.9    |
| SBC1    | Southern Branch          | Pollution Sensitive Species Abundance  | 0.0500  | 39.87         | 14.61   | 73.1     |
| SBD2    | Southern Branch          | Pollution Sensitive Species Biomass    | 0.0500  | 26.14         | 11.91   | 59.6     |
| SBD4    | Southern Branch          | Shannon-Weiner Diversity Index         | 0.0500  | 2.41          | -0.25   | -51.1    |
| WBB1    | Western Branch           | Pollution Sensitive Species Abundance  | 0.0143  | 52.08         | 8.14    | 40.7     |
| WBB1    | Western Branch           | Pollution Indicative Species Abundance | 0.0500  | 35.74         | -7.11   | -35.6    |
| WBB1    | Western Branch           | Pollution Sensitive Species Biomass    | 0.0500  | 13.60         | 10.62   | 53.1     |
| WBB5    | Western Branch           | Pollution Sensitive Species Abundance  | 0.0500  | 24.58         | 17.09   | 85.5     |

Table 9. Comparison of nutrient concentrations in the Elizabeth River compare to the lower polyhaline section of the James River (JMSPH see Appendix B, Fig. B1). All values are in mg/l except chlorophyl *a* (CHLA) which is in µg/l and are the median values for the last three years of collection. Data for JMSPH from Dauer et al. 2005.

| Parameter | James River (JMSPH) | Elizabeth River Mainstem | Western Branch | Eastern Branch | Southern Branch |
|-----------|---------------------|--------------------------|----------------|----------------|-----------------|
| STN       | 0.46                | 0.65                     | 0.70           | 0.84           | 1.12            |
| SDIN      | 0.06                | 0.28                     | 0.20           | 0.40           | 0.58            |
| STP       | 0.04                | 0.05                     | 0.06           | 0.05           | 0.06            |
| SDIP      | 0.01                | 0.02                     | 0.02           | 0.02           | 0.03            |
| CHLA      | 7.57                | 10.23                    | 11.64          | 6.09           | 3.62            |

**Appendix A:** Metrics and thresholds for calculating the Benthic Index Biotic Integrity

Table A1. Thresholds used to score each metric of the Chesapeake Bay B-IBI. Updated for the tidal freshwater and oligohaline habitats, and corrected from Weisberg et al. (1997) for the high mesohaline mud and polyhaline sand habitats.

| Scoring Criteria                               |             |                            |                |
|--|-------------|----------------------------|----------------|
|  | 5           | 3                          | 1              |
| <b>Tidal Freshwater</b>                        |             |                            |                |
| Abundance (#/m <sup>2</sup> )                  | ≥ 1050-4000 | 800-1050 or<br>≥ 4000-5500 | <800 or ≥ 5500 |
| Abundance of pollution-indicative taxa (%)     | ≤ 39        | 39-87                      | >87            |
| Abundance of deep-deposit feeders (%)          | ≤ 70        | 70-95                      | >95            |
| Tolerance Score                                | ≤ 8         | 8-9.35                     | >9.35          |
| <b>Oligohaline</b>                             |             |                            |                |
| Abundance (#/m <sup>2</sup> )                  | ≥ 450-3350  | 180-450 or<br>≥ 3350-4050  | <180 or ≥ 4050 |
| Abundance of pollution-indicative taxa (%)     | ≤ 27        | 27-95                      | >95            |
| Abundance of pollution-sensitive taxa (%)      | ≥ 26        | 0.2-26                     | <0.2           |
| Abundance of carnivores and omnivores (%)      | ≥ 35        | 15-35                      | <15            |
| Tolerance Score                                | ≤ 6         | 6-9.05                     | >9.05          |
| Tanypodini to Chironomidae abundance ratio (%) | ≤ 17        | 17-64                      | >64            |
| <b>Low Mesohaline</b>                          |             |                            |                |
| Shannon-Wiener                                 | ≥ 2.5       | 1.7-2.5                    | <1.7           |
| Abundance (#/m <sup>2</sup> )                  | ≥ 1500-2500 | 500-1500 or<br>≥ 2500-6000 | <500 or ≥ 6000 |
| Biomass (g/m <sup>2</sup> )                    | ≥ 5-10      | 1-5 or ≥ 10-30             | <1 or ≥ 30     |
| Abundance of pollution-indicative taxa (%)     | ≤ 10        | 10-20                      | >20            |
| Biomass of pollution-sensitive taxa (%)        | ≥ 80        | 40-80                      | <40            |
| Biomass deeper than 5 cm (%)                   | ≥ 80        | 10-80                      | <10            |



Table A1. Continued.

| Scoring Criteria                           |             |                             |                 |
|--|-------------|-----------------------------|-----------------|
|  | 5           | 3                           | 1               |
| <b>High Mesohaline Sand</b>                |             |                             |                 |
| Shannon-Wiener                             | ≥ 3.2       | 2.5-3.2                     | <2.5            |
| Abundance (#/m <sup>2</sup> )              | ≥ 1500-3000 | 1000-1500 or<br>≥ 3000-5000 | <1000 or ≥ 5000 |
| Biomass (g/m <sup>2</sup> )                | ≥ 3-15      | 1-3 or ≥ 15-50              | <1 or ≥ 50      |
| Abundance of pollution-indicative taxa (%) | ≤ 10        | 10-25                       | >25             |
| Abundance of pollution-sensitive taxa (%)  | ≥ 40        | 10-40                       | <10             |
| Abundance of carnivores and omnivores (%)  | ≥ 35        | 20-35                       | <20             |
| <b>High Mesohaline Mud</b>                 |             |                             |                 |
| Shannon-Wiener                             | 3.0         | 2.0-3.0                     | <2.0            |
| Abundance (#/m <sup>2</sup> )              | ≥ 1500-2500 | 1000-1500 or<br>≥ 2500-5000 | <1000 or ≥ 5000 |
| Biomass (g/m <sup>2</sup> )                | ≥ 2-10      | 0.5-2 or ≥ 10-50            | <0.5 or ≥ 50    |
| Biomass of pollution-indicative taxa (%)   | ≤ 5         | 5-30                        | >30             |
| Biomass of pollution-sensitive taxa (%)    | ≥ 60        | 30-60                       | <30             |
| Abundance of carnivores and omnivores (%)  | ≥ 25        | 10-25                       | <10             |
| Biomass deeper than 5 cm (%)               | ≥ 60        | 10-60                       | <10             |
| <b>Polyhaline Sand</b>                     |             |                             |                 |
| Shannon-Wiener                             | ≥ 3.5       | 2.7-3.5                     | <2.7            |
| Abundance (#/m <sup>2</sup> )              | ≥ 3000-5000 | 1500-3000 or<br>≥ 5000-8000 | <1500 or ≥ 8000 |
| Biomass (g/m <sup>2</sup> )                | ≥ 5-20      | 1-5 or ≥ 20-50              | <1 or ≥ 50      |
| Biomass of pollution-indicative taxa (%)   | ≤ 5         | 5-15                        | >15             |
| Abundance of pollution-sensitive taxa (%)  | ≥ 50        | 25-50                       | <25             |
| Abundance of deep-deposit feeders (%)      | ≥ 25        | 10-25                       | <10             |


Table 1. Continued.

| Scoring Criteria   |                   |                                   |                         |
|--|-------------------|-----------------------------------|-------------------------|
|  | 5                 | 3                                 | 1                       |
| <b>Polyhaline Mud</b>  |                   |                                   |                         |
| Shannon-Wiener   | $\geq 3.3$        | 2.4-3.3                           | $< 2.4$                 |
| Abundance ( $\#/m^2$ )   | $\geq 1500$ -3000 | 1000-1500 or<br>$\geq 3000$ -8000 | $< 1000$ or $\geq 8000$ |
| Biomass ( $g/m^2$ )  | $\geq 3$ -10      | 0.5-3 or $\geq 10$ -30            | $< 0.5$ or $\geq 30$    |
| Biomass of pollution-indicative taxa (%)                       | $\leq 5$          | 5-20                              | $> 20$                  |
| Biomass of pollution-sensitive taxa (%)                        | $\geq 60$         | 30-60                             | $< 30$                  |
| Abundance of carnivores and omnivores                          | $\geq 40$         | 25-40                             | $< 25$                  |
| Number of taxa $> 5$ cm below the sediment-water interface (%) | $\geq 40$         | 10-40                             | $< 10$                  |


## **Appendix B:** Summary of water quality status and trends for the Elizabeth River

### **Preface:**

In this appendix water quality status and trends for the Elizabeth River are summarized. These data are collected by the Virginia Water Quality Monitoring Program as part of the Chesapeake Bay Restoration Program. Details of collection and laboratory methodology can be found in Dauer et al. 2005 which can be downloaded in pdf format from the Old Dominion University Chesapeake Bay Program website <[www.chesapeakebay.odu.edu](http://www.chesapeakebay.odu.edu)> under “Reports.” The James River Report includes the Elizabeth River, the Chickahominy River and the Appomattox River. The York River Report includes the tidal Pamunkey River and Mattaponi River. The Rappahannock River Report includes the Corrotoman River. Also available at this website are appendices that include (1) tables of status for all parameters measured at all stations sampled by each program, (2) tables of all parameters and metrics for which there was a significant trend, and (3) scatter plots of all parameters over time. There are five appendices: water quality, phytoplankton, primary productivity, zooplankton and benthos.



**Figure B1.** Map showing the locations of the water quality monitoring stations in the Virginia tributaries and the Lower Chesapeake Bay Mainstem (Dauer et al 2005). Insert shows location of Elizabeth River monitoring stations. Also shown are ellipses that delineate the Chesapeake Bay Program segmentation scheme.



**Figure B2.** Map of the Elizabeth River basin showing summaries of the status and trend analyses for each segment. Abbreviations: TN - total nitrogen; DIN - dissolved inorganic nitrogen; TP - total phosphorus; DIP - dissolved inorganic phosphorus. The prefixes S and B refer to surface and bottom measurements.




Figure B3. Map of the Elizabeth River basin showing summaries of the status and trend analyses for each segment. Abbreviations: SCHLA -surface chlorophyll a; TSS- total suspended solids, SECCHI - secchi depth, BDO - bottom dissolved oxygen; WTEMP - water temperature, SALIN - salinity. The prefixes S and B refer to surface and bottom measurements.

Table B-1. Status in water quality environmental indicators in the Elizabeth River. Status designations determined by the Chesapeake Bay Program for the three year period 2002 through 2004. For information about field collection, laboratory analyses and status determination see Dauer et al. 2005. Secchi depth in meters, chlorophyll *a* in µg/l, all others in mg/l. S is surface and B is bottom layer. WBEMH - Western Branch, SBEMH - Southern Branch, EBEMH - Eastern Branch, ELIPH - Elizabeth River Mainstem.

| Segment | Season | Parameter | Median | Score | Status |
|---------|--------|-----------|--------|-------|--------|
| EBEMH   | STN    | Annual    | 0.845  | 80.5  | Poor   |
| EBEMH   | BTN    | Annual    | 0.768  | 76.0  | Poor   |
| EBEMH   | SDIN   | Annual    | 0.395  | 88.4  | Poor   |
| EBEMH   | BDIN   | Annual    | 0.346  | 93.8  | Poor   |
| EBEMH   | STP    | Annual    | 0.053  | 72.3  | Poor   |
| EBEMH   | BTP    | Annual    | 0.052  | 52.6  | Fair   |
| EBEMH   | SPO4F  | Annual    | 0.021  | 84.7  | Poor   |
| EBEMH   | BPO4F  | Annual    | 0.028  | 84.2  | Poor   |
| EBEMH   | SCHLA  | Annual    | 6.09   | 29.2  | Good   |
| EBEMH   | STSS   | Annual    | 8.31   | 42.5  | Fair   |
| EBEMH   | BTSS   | Annual    | 11.36  | 29.2  | Good   |
| EBEMH   | SECCHI | Annual    | 1.00   | 25.3  | Poor   |
| WBEMH   | STN    | Annual    | 0.698  | 57.4  | Fair   |
| WBEMH   | BTN    | Annual    | 0.691  | 68.1  | Poor   |
| WBEMH   | SDIN   | Annual    | 0.199  | 59.4  | Fair   |
| WBEMH   | BDIN   | Annual    | 0.214  | 68.8  | Poor   |
| WBEMH   | STP    | Annual    | 0.059  | 70.0  | Poor   |
| WBEMH   | BTP    | Annual    | 0.058  | 54.3  | Fair   |
| WBEMH   | SPO4F  | Annual    | 0.017  | 74.8  | Poor   |
| WBEMH   | BPO4F  | Annual    | 0.019  | 63.5  | Poor   |
| WBEMH   | SCHLA  | Annual    | 11.64  | 56.8  | Fair   |
| WBEMH   | STSS   | Annual    | 15.93  | 79.1  | Poor   |
| WBEMH   | BTSS   | Annual    | 20.21  | 65.1  | Poor   |
| WBEMH   | SECCHI | Annual    | 0.70   | 9.8   | Poor   |
| SBEMH   | STN    | Annual    | 1.115  | 95.6  | Poor   |
| SBEMH   | BTN    | Annual    | 0.989  | 91.6  | Poor   |
| SBEMH   | SDIN   | Annual    | 0.584  | 96.1  | Poor   |
| SBEMH   | BDIN   | Annual    | 0.497  | 98.2  | Poor   |
| SBEMH   | STP    | Annual    | 0.060  | 65.0  | Fair   |
| SBEMH   | BTP    | Annual    | 0.062  | 64.6  | Poor   |
| SBEMH   | SPO4F  | Annual    | 0.031  | 96.2  | Poor   |
| SBEMH   | BPO4F  | Annual    | 0.035  | 92.4  | Poor   |
| SBEMH   | SCHLA  | Annual    | 3.62   | 7.9   | Good   |
| SBEMH   | STSS   | Annual    | 7.75   | 35.6  | Good   |
| SBEMH   | BTSS   | Annual    | 9.28   | 24.3  | Good   |
| SBEMH   | SECCHI | Annual    | 0.90   | 16.0  | Poor   |
| ELIPH   | STN    | Annual    | 0.645  | 55.2  | Fair   |
| ELIPH   | BTN    | Annual    | 0.603  | 64.0  | Fair   |
| ELIPH   | SDIN   | Annual    | 0.277  | 72.6  | Poor   |
| ELIPH   | BDIN   | Annual    | 0.225  | 87.5  | Poor   |
| ELIPH   | STP    | Annual    | 0.054  | 59.2  | Fair   |
| ELIPH   | BTP    | Annual    | 0.063  | 72.6  | Poor   |
| ELIPH   | SPO4F  | Annual    | 0.021  | 85.2  | Poor   |
| ELIPH   | BPO4F  | Annual    | 0.024  | 77.1  | Poor   |
| ELIPH   | SCHLA  | Annual    | 10.23  | 58.4  | Fair   |
| ELIPH   | STSS   | Annual    | 10.31  | 54.3  | Fair   |
| ELIPH   | BTSS   | Annual    | 18.02  | 48.0  | Fair   |
| ELIPH   | SECCHI | Annual    | 0.95   | 19.1  | Poor   |

Table B-2. Blocked seasonal Kendall Long-term trends in water quality for the segment ELIPH in the Elizabeth River (1985-2004). Parameters are as follows: STN=Surface total nitrogen, BTN=Bottom total nitrogen, SDIN=Surface dissolved inorganic nitrogen, BDIN=Bottom dissolved inorganic nitrogen, STP=Surface total phosphorus, BTP=Bottom total phosphorus, SPO4F=Surface dissolved inorganic phosphorus, BPO4F=Bottom dissolved inorganic phosphorus.


| Segment | Parameter | '85-'93<br>Trend<br>p value | '85-93<br>Trend<br>slope | '85-'93<br>Trend<br>Direction | '95-'02<br>Trend<br>p value | '95-'02<br>Trend<br>Slope | '95-'02<br>Trend<br>Direction | Trend<br>Comparison<br>p value | Trend<br>Comparison<br>Significance | Combined<br>Trend<br>p value | Combined<br>Trend<br>Direction |
|---------|-----------|-----------------------------|--------------------------|-------------------------------|-----------------------------|---------------------------|-------------------------------|--------------------------------|-------------------------------------|------------------------------|--------------------------------|
| ELIPH   | STN       | 1.0000                      | 0.000                    | No Trend                      | 0.1000                      | 0.006                     | No Trend                      | 0.1761                         | Same                                | 0.1900                       | No Trend                       |
| ELIPH   | BTN       | 1.0000                      | 0.000                    | No Trend                      | 0.0201                      | 0.007                     | No Trend                      | 0.0570                         | Same                                | 0.0570                       | No Trend                       |
| ELIPH   | SDIN      | 0.9426                      | 0.000                    | No Trend                      | 0.0072                      | 0.009                     | Degrading                     | 0.0341                         | Same                                | 0.0248                       | No Trend                       |
| ELIPH   | BDIN      | 0.8504                      | 0.000                    | No Trend                      | 0.0389                      | 0.005                     | No Trend                      | 0.1149                         | Same                                | 0.0669                       | No Trend                       |
| ELIPH   | STP       | 0.8184                      | 0.000                    | No Trend                      | 0.0663                      | -0.001                    | No Trend                      | 0.1087                         | Same                                | 0.2008                       | No Trend                       |
| ELIPH   | BTP       | 0.1060                      | 0.001                    | No Trend                      | 0.0707                      | -0.001                    | No Trend                      | 0.0146                         | Same                                | 0.7015                       | No Trend                       |
| ELIPH   | SPO4F     | 0.7632                      | 0.000                    | No Trend                      | 0.8765                      | 0.000                     | No Trend                      | 0.7431                         | Same                                | 0.9347                       | No Trend                       |
| ELIPH   | SPO4F     | 0.5036                      | 0.000                    | No Trend                      | 0.7564                      | 0.000                     | No Trend                      | 0.8540                         | Same                                | 0.4998                       | No Trend                       |



**Appendix C:** Summary of benthic community status and trends for the Elizabeth River stations (SBE2 and SBE5) of the Benthic Monitoring Program of the Chesapeake Bay Program.

**Preface:**

In this appendix status and trends in the benthic community for the Elizabeth River are summarized. These data are collected by the Virginia Benthic Monitoring Program as part of the Chesapeake Bay Restoration Program. Details of collection and laboratory methodology can be found in Dauer et al. 2005 which can be downloaded in pdf format from the Old Dominion University Chesapeake Bay Program website <[www.chesapeakebay.odu.edu](http://www.chesapeakebay.odu.edu)> under “Reports.” This appendix presents a summary of status and trends for stations SBE2 and SBE5 which are located in the Southern Branch of the river. Shown are scatter plots of the B-IBI and for several benthic metrics for the period 1989 to 2004.



**Figure C1.** Location of living resource monitoring stations in the Virginia tributaries and the Lower Chesapeake Bay Mainstem. Insert shows benthic monitoring stations in the Southern Branch (SBE2 and SBE5) established in 1989.




Figure C2. Map of the James River basin showing summaries of the status and trend analyses for the B-IBI and selected metrics for each segment for the period 1985 through 2004. Stations SBE2 and SBE5 are in the Elizabeth River.

Table C1. Annual season term trends in the benthic IBI and its component metrics in the James River and Elizabeth River for the period of 1985 through 2004.

| Station | Parameter                              | P value | Slope  | Baseline | % Change | Direction |
|---------|--|---------|--------|----------|----------|-----------|
| TF5.5   | Benthic Index of Biotic Integrity      | 0.1598  | 0.04   | 2.13     | 40.06    | No Trend  |
| TF5.5   | Total Abundance per square meter       | 0.1534  | 208.69 | 1335.60  | 312.50   | No Trend  |
| TF5.5   | Total Biomass per square meter         | 0.4168  | 0.03   | 0.34     | 159.61   | No Trend  |
| TF5.5   | Shannon-Weiner Diversity Index         | 0.8457  | 0.00   | 1.18     | 7.66     | No Trend  |
| TF5.5   | Pollution Sensitive Species Abundance  | 0.0042  | 1.33   | 0.00     | N.E.     | Improving |
| TF5.5   | Pollution Indicative Species Abundance | 0.3636  | -0.46  | 20.14    | -45.27   | No Trend  |
| TF5.5   | Pollution Sensitive Species Biomass    | 0.0042  | 3.01   | 0.00     | N.E.     | Improving |
| TF5.5   | Pollution Indicative Species Biomass   | 0.2992  | -1.38  | 43.89    | -63.00   | No Trend  |
| RET5.2  | Benthic Index of Biotic Integrity      | 0.0013  | 0.06   | 1.92     | 63.24    | Improving |
| RET5.2  | Total Abundance per square meter       | 0.2055  | 27.30  | 610.56   | 89.41    | No Trend  |
| RET5.2  | Total Biomass per square meter         | 0.7208  | -0.01  | 7.34     | -1.84    | No Trend  |
| RET5.2  | Shannon-Weiner Diversity Index         | 0.0137  | 0.05   | 1.66     | 56.55    | Improving |
| RET5.2  | Pollution Sensitive Species Abundance  | 0.0012  | 1.58   | 10.42    | 304.13   | Improving |
| RET5.2  | Pollution Indicative Species Abundance | 0.8419  | 0.00   | 16.67    | 0.00     | No Trend  |
| RET5.2  | Pollution Sensitive Species Biomass    | 0.0137  | 2.23   | 28.04    | 158.94   | Improving |
| RET5.2  | Pollution Indicative Species Biomass   | 0.3352  | -0.01  | 8.42     | -1.67    | No Trend  |
| LE5.1   | Benthic Index of Biotic Integrity      | 0.3866  | 0.02   | 3.02     | 12.52    | No Trend  |
| LE5.1   | Total Abundance per square meter       | 0.8691  | 6.13   | 524.70   | 19.87    | No Trend  |
| LE5.1   | Total Biomass per square meter         | 0.3648  | -0.03  | 5.71     | -10.01   | No Trend  |
| LE5.1   | Shannon-Weiner Diversity Index         | 0.8048  | 0.01   | 2.06     | 5.08     | No Trend  |
| LE5.1   | Pollution Sensitive Species Abundance  | 0.5641  | -0.29  | 32.16    | -15.32   | No Trend  |
| LE5.1   | Pollution Indicative Species Abundance | 0.1554  | 0.28   | 7.13     | 66.00    | No Trend  |
| LE5.1   | Pollution Sensitive Species Biomass    | 0.8691  | -0.22  | 74.24    | -4.96    | No Trend  |
| LE5.1   | Pollution Indicative Species Biomass   | 0.1554  | 0.12   | 4.82     | 42.11    | No Trend  |
| LE5.2   | Benthic Index of Biotic Integrity      | 0.5811  | -0.02  | 3.33     | -10.94   | No Trend  |
| LE5.2   | Total Abundance per square meter       | 0.0976  | 41.62  | 1221.12  | 68.17    | Improving |
| LE5.2   | Total Biomass per square meter         | 0.3978  | -0.12  | 4.58     | -54.17   | No Trend  |
| LE5.2   | Shannon-Weiner Diversity Index         | 0.2992  | -0.03  | 2.55     | -21.36   | No Trend  |
| LE5.2   | Pollution Sensitive Species Abundance  | 0.0443  | 2.07   | 34.48    | 119.99   | Improving |
| LE5.2   | Pollution Indicative Species Abundance | 0.9483  | -0.07  | 17.25    | -8.48    | No Trend  |
| LE5.2   | Pollution Sensitive Species Biomass    | 0.7952  | 0.66   | 40.78    | 32.52    | No Trend  |
| LE5.2   | Pollution Indicative Species Biomass   | 0.1194  | 0.38   | 8.76     | 87.02    | No Trend  |
| LE5.4   | Benthic Index of Biotic Integrity      | 0.8704  | 0.00   | 3.73     | 0.00     | No Trend  |
| LE5.4   | Total Abundance per square meter       | 0.5164  | 19.95  | 2528.10  | 15.78    | No Trend  |
| LE5.4   | Total Biomass per square meter         | 0.0023  | -1.54  | 22.86    | -134.39  | Degrading |
| LE5.4   | Shannon-Weiner Diversity Index         | 0.0798  | -0.02  | 3.69     | -12.86   | Degrading |
| LE5.4   | Pollution Sensitive Species Abundance  | 0.2176  | -0.71  | 50.52    | -27.97   | No Trend  |
| LE5.4   | Pollution Indicative Species Abundance | 0.1214  | -0.07  | 1.99     | -71.06   | No Trend  |
| LE5.4   | Pollution Sensitive Species Biomass    | 0.0231  | -1.72  | 60.10    | -57.39   | Degrading |
| LE5.4   | Pollution Indicative Species Biomass   | 0.9737  | 0.00   | 0.51     | 0.00     | No Trend  |
| SBE2    | Benthic Index of Biotic Integrity      | 0.2403  | 0.02   | 2.00     | 19.37    | No Trend  |
| SBE2    | Total Abundance per square meter       | 0.0271  | 186.12 | 1659.96  | 179.39   | Improving |
| SBE2    | Total Biomass per square meter         | 0.3214  | 0.03   | 0.92     | 49.07    | No Trend  |
| SBE2    | Shannon-Weiner Diversity Index         | 0.4713  | 0.02   | 1.67     | 20.62    | No Trend  |
| SBE2    | Pollution Sensitive Species Abundance  | 0.0003  | 4.77   | 4.62     | 1649.97  | Improving |
| SBE2    | Pollution Indicative Species Abundance | 0.0192  | -3.12  | 72.25    | -69.19   | Improving |
| SBE2    | Pollution Sensitive Species Biomass    | 0.0272  | 2.16   | 8.41     | 411.33   | Improving |
| SBE2    | Pollution Indicative Species Biomass   | 0.8571  | -0.15  | 44.94    | -5.17    | No Trend  |
| SBE5    | Benthic Index of Biotic Integrity      | 0.0050  | 0.07   | 1.25     | 95.59    | Improving |
| SBE5    | Total Abundance per square meter       | 0.7187  | 50.56  | 3806.46  | 21.25    | No Trend  |
| SBE5    | Total Biomass per square meter         | 0.1612  | 0.05   | 0.58     | 131.15   | No Trend  |
| SBE5    | Shannon-Weiner Diversity Index         | 0.5890  | 0.02   | 0.99     | 36.55    | No Trend  |
| SBE5    | Pollution Sensitive Species Abundance  | 0.0000  | 4.01   | 0.31     | 20503.15 | Improving |
| SBE5    | Pollution Indicative Species Abundance | 0.0012  | -5.13  | 85.43    | -96.15   | Improving |
| SBE5    | Pollution Sensitive Species Biomass    | 0.0060  | 2.79   | 2.15     | 2072.05  | Improving |
| SBE5    | Pollution Indicative Species Biomass   | 0.0717  | -3.16  | 77.83    | -65.06   | Improving |

## SBE2 (2-SB)




Figure C-3. Plot of the benthic IBI against time at station SBE2 for 1985 through 2004.

## SBE5 (5-SB)




Figure C-4. Plot of the benthic IBI against time at station SBE5 for 1985 through 2004.

## SBE2 (2—SB)




Figure C-5. Plot of total benthic community abundance at station SBE2 for 1985 through 2004.

## SBE5 (5—SB)




Figure C-6. Plot of total benthic community abundance at station SBE5 for 1985 through 2004.

## SBE2 (2—SB)




Figure C-7. Plot of total benthic community biomass at station SBE2 for 1985 through 2004.

## SBE5 (5—SB)




Figure C-8. Plot of total benthic community biomass at station SBE5 for 1985 through 2004.

## SBE2 (2—SB)




Figure C-9. Plot of pollution sensitive species abundance at station SBE2 for 1985 through 2004.

## SBE5 (5—SB)




Figure C-10. Plot of pollution sensitive species abundance at station SBE5 for 1985 through 2004.



## SBE2 (2—SB)




Figure C-11. Plot of pollution indicative species abundance at station SBE2 for 1985 through 2004.

## SBE5 (5—SB)




Figure C-12. Plot of pollution indicative species abundance at station SBE5 for 1985 through 2004.

## SBE2 (2—SB)




Figure C-13. Plot of pollution sensitive species biomass at station SBE2 for 1985 through 2004.

## SBE5 (5—SB)




Figure C-14. Plot of pollution sensitive species biomass at station SBE5 for 1985 through 2004.

## SBE2 (2-SB)




Figure C-15. Plot of pollution indicative species biomass at station SBE2 for 1985 through 2004.

## SBE5 (5-SB)




Figure C-16. Plot of pollution indicative species biomass at station SBE5 for 1985 through 2004.

## SBE2 (2-SB)




Figure C-17. Plot of Shannon-Weiner diversity index at station SBE2 for 1985 through 2004.

## SBE5 (5-SB)




Figure C-18. Plot of Shannon-Weiner diversity index at station SBE5 for 1985 through 2004.

**Appendix D:** Plots of fixed -point station B-IBI and metric values from 1999-2004 for the 14 stations of the Elizabeth River Benthic Monitoring Program

|               |   |       |
|---------------|---|-------|
| Figure D-1.   | Plot of the benthic IBI at station ELC1 from 1999 through 2004. ....                      | D-59  |
| Figure D- 2.  | Plot of total benthic community abundance at station ELC1 for 1999 through 2004. ...      | D-60  |
| Figure D- 3.  | Plot of total benthic community biomass at station ELC1 for 1999 through 2004. ....       | D-61  |
| Figure D- 4.  | Plot of the Shannon-Weiner diversity index at station ELC1 for 1999 through 2004. ...     | D-62  |
| Figure D- 5.  | Plot of pollution sensitive species abundance at station ELC1for 1999 through 2004. ...   | D-63  |
| Figure D- 6.  | Plot of pollution indicative species abundance at station ELC1 for 1999 through 2004. ... | D-64  |
| Figure D- 7.  | Plot of pollution sensitive species biomass at station ELC1 for 1999 through 2004. ...    | D-65  |
| Figure D- 8.  | Plot of pollution indicative species biomass at station ELC1 for 1999 through 2004. ...   | D-66  |
| Figure D- 9.  | Plot of the benthic IBI at station ELD1 from 1999 through 2004. ....                      | D-67  |
| Figure D- 10. | Plot of total benthic community abundance at station ELD1 for 1999 through 2004. ...      | D-68  |
| Figure D- 11. | Plot of total benthic community biomass at station ELD1 for 1999 through 2004. ....       | D-69  |
| Figure D- 12. | Plot of the Shannon-Weiner diversity index at station ELD1 for 1999 through 2004. ...     | D-70  |
| Figure D- 13. | Plot of pollution sensitive species abundance at station ELD1for 1999 through 2004. ...   | D-71  |
| Figure D- 14. | Plot of pollution indicative species abundance at station ELD1 for 1999 through 2004. ... | D-72  |
| Figure D- 15. | Plot of pollution sensitive species biomass at station ELD1 for 1999 through 2004. ...    | D-73  |
| Figure D- 16. | Plot of pollution indicative species biomass at station ELD1 for 1999 through 2004. ...   | D-74  |
| Figure D- 17. | Plot of the benthic IBI at station ELF1 from 1999 through 2004. ....                      | D-75  |
| Figure D- 18. | Plot of total benthic community abundance at station ELF1 for 1999 through 2004. ...      | D-76  |
| Figure D- 19. | Plot of total benthic community biomass at station ELF1 for 1999 through 2004. ....       | D-77  |
| Figure D- 20. | Plot of the Shannon-Weiner diversity index at station ELF1 for 1999 through 2004. ...     | D-78  |
| Figure D- 21. | Plot of pollution sensitive species abundance at station ELF1for 1999 through 2004. ...   | D-79  |
| Figure D- 22. | Plot of pollution indicative species abundance at station ELF1 for 1999 through 2004. ... | D-80  |
| Figure D- 23. | Plot of pollution sensitive species biomass at station ELF1 for 1999 through 2004. ...    | D-81  |
| Figure D- 24. | Plot of pollution indicative species biomass at station ELF1 for 1999 through 2004. ...   | D-82  |
| Figure D- 25. | Plot of the benthic IBI at station EBB1 from 1999 through 2004. ....                      | D-83  |
| Figure D- 26. | Plot of total benthic community abundance at station EBB1 for 1999 through 2004. ...      | D-84  |
| Figure D- 27. | Plot of total benthic community biomass at station EBB1 for 1999 through 2004. ....       | D-85  |
| Figure D- 28. | Plot of the Shannon-Weiner diversity index at station EBB1 for 1999 through 2004. ...     | D-86  |
| Figure D- 29. | Plot of pollution sensitive species abundance at station EBB1for 1999 through 2004. ...   | D-87  |
| Figure D- 30. | Plot of pollution indicative species abundance at station EBB1 for 1999 through 2004. ... | D-88  |
| Figure D- 31. | Plot of pollution sensitive species biomass at station EBB1 for 1999 through 2004. ...    | D-89  |
| Figure D- 32. | Plot of pollution indicative species biomass at station EBB1 for 1999 through 2004. ...   | D-90  |
| Figure D- 33. | Plot of the benthic IBI at station LFA1 from 1999 through 2004. ....                      | D-91  |
| Figure D- 34. | Plot of total benthic community abundance at station LFA1 for 1999 through 2004. ...      | D-92  |
| Figure D- 35. | Plot of total benthic community biomass at station LFA1 for 1999 through 2004. ....       | D-93  |
| Figure D- 36. | Plot of the Shannon-Weiner diversity index at station LFA1 for 1999 through 2004. ...     | D-94  |
| Figure D- 37. | Plot of pollution sensitive species abundance at station LFA1for 1999 through 2004. ...   | D-95  |
| Figure D- 38. | Plot of pollution indicative species abundance at station LFA1 for 1999 through 2004. ... | D-96  |
| Figure D- 39. | Plot of pollution sensitive species biomass at station LFA1 for 1999 through 2004. ...    | D-97  |
| Figure D- 40. | Plot of pollution indicative species biomass at station LFA1 for 1999 through 2004. ...   | D-98  |
| Figure D- 41. | Plot of the benthic IBI at station LFB1 from 1999 through 2004. ....                      | D-99  |
| Figure D- 42. | Plot of total benthic community abundance at station LFB1 for 1999 through 2004. ...      | D-100 |
| Figure D- 43. | Plot of total benthic community biomass at station LFB1 for 1999 through 2004. ....       | D-101 |
| Figure D- 44. | Plot of the Shannon-Weiner diversity index at station LFB1 for 1999 through 2004. ...     | D-102 |
| Figure D- 45. | Plot of pollution sensitive species abundance at station LFB1for 1999 through 2004. ...   | D-103 |
| Figure D- 46. | Plot of pollution indicative species abundance at station LFB1 for 1999 through 2004. ... | D-104 |
| Figure D- 47. | Plot of pollution sensitive species biomass at station LFB1 for 1999 through 2004. ...    | D-105 |
| Figure D- 48. | Plot of pollution indicative species biomass at station LFB1 for 1999 through 2004. ...   | D-106 |
| Figure D- 49. | Plot of the benthic IBI at station WBB1 from 1999 through 2004. ....                      | D-107 |
| Figure D- 50. | Plot of total benthic community abundance at station WBB1 for 1999 through 2004. ...      | D-108 |
| Figure D- 51. | Plot of total benthic community biomass at station WBB1 for 1999 through 2004. ...        | D-109 |
| Figure D- 52. | Plot of the Shannon-Weiner diversity index at station WBB1 for 1999 through 2004. ...     | D-110 |
| Figure D- 53. | Plot of pollution sensitive species abundance at station WBB1for 1999 through 2004. ...   | D-111 |
| Figure D- 54. | Plot of pollution indicative species abundance at station WBB1 for 1999 through 2004. ... | D-112 |

|                |  |       |
|----------------|--|-------|
| Figure D- 55.  | Plot of pollution sensitive species biomass at station WBB1 for 1999 through 2004. .     | D-113 |
| Figure D- 56.  | Plot of pollution indicative species biomass at station WBB1 for 1999 through 2004. .    | D-114 |
| Figure D- 57.  | Plot of the benthic IBI at station WBB5 from 1999 through 2004. ....                     | D-115 |
| Figure D- 58.  | Plot of total benthic community abundance at station WBB5 for 1999 through 2004. .       | D-116 |
| Figure D- 59.  | Plot of total benthic community biomass at station WBB5 for 1999 through 2004. . .       | D-117 |
| Figure D- 60.  | Plot of the Shannon-Weiner diversity index at station WBB5 for 1999 through 2004. .      | D-118 |
| Figure D- 61.  | Plot of pollution sensitive species abundance at station WBB5for 1999 through 2004. .    | D-119 |
| Figure D- 62.  | Plot of pollution indicative species abundance at station WBB5 for 1999 through 2004.D-  | D-120 |
| Figure D- 63.  | Plot of pollution sensitive species biomass at station WBB5 for 1999 through 2004. .     | D-121 |
| Figure D- 64.  | Plot of pollution indicative species biomass at station WBB5 for 1999 through 2004. .    | D-122 |
| Figure D- 65.  | Plot of the benthic IBI at station SBA1 from 1999 through 2004. ....                     | D-123 |
| Figure D- 66.  | Plot of total benthic community abundance at station SBA1 for 1999 through 2004. . .     | D-124 |
| Figure D- 67.  | Plot of total benthic community biomass at station SBA1 for 1999 through 2004. . . .     | D-125 |
| Figure D- 68.  | Plot of the Shannon-Weiner diversity index at station SBA1 for 1999 through 2004. .      | D-126 |
| Figure D- 69.  | Plot of pollution sensitive species abundance at station SBA1for 1999 through 2004. .    | D-127 |
| Figure D- 70.  | Plot of pollution indicative species abundance at station SBA1 for 1999 through 2004. D- | D-128 |
| Figure D- 71.  | Plot of pollution sensitive species biomass at station SBA1 for 1999 through 2004. . .   | D-129 |
| Figure D- 72.  | Plot of pollution indicative species biomass at station SBA1 for 1999 through 2004. .    | D-130 |
| Figure D- 73.  | Plot of the benthic IBI at station SBB1 from 1999 through 2004. ....                     | D-131 |
| Figure D- 74.  | Plot of total benthic community abundance at station SBB1 for 1999 through 2004. . .     | D-132 |
| Figure D- 75.  | Plot of total benthic community biomass at station SBB1 for 1999 through 2004. . . .     | D-133 |
| Figure D- 76.  | Plot of the Shannon-Weiner diversity index at station SBB1 for 1999 through 2004. .      | D-134 |
| Figure D- 77.  | Plot of pollution sensitive species abundance at station SBB1for 1999 through 2004. .    | D-135 |
| Figure D- 78.  | Plot of pollution indicative species abundance at station SBB1 for 1999 through 2004. D- | D-136 |
| Figure D- 79.  | Plot of pollution sensitive species biomass at station SBB1 for 1999 through 2004. . .   | D-137 |
| Figure D- 80.  | Plot of pollution indicative species biomass at station SBB1 for 1999 through 2004. .    | D-138 |
| Figure D- 81.  | Plot of the benthic IBI at station SBC1 from 1999 through 2004. ....                     | D-139 |
| Figure D- 82.  | Plot of total benthic community abundance at station SBC1 for 1999 through 2004. . .     | D-140 |
| Figure D- 83.  | Plot of total benthic community biomass at station SBC1 for 1999 through 2004. . . .     | D-141 |
| Figure D- 84.  | Plot of the Shannon-Weiner diversity index at station SBC1 for 1999 through 2004. .      | D-142 |
| Figure D- 85.  | Plot of pollution sensitive species abundance at station SBC1for 1999 through 2004. .    | D-143 |
| Figure D- 86.  | Plot of pollution indicative species abundance at station SBC1 for 1999 through 2004. D- | D-144 |
| Figure D- 87.  | Plot of pollution sensitive species biomass at station SBC1 for 1999 through 2004. . .   | D-145 |
| Figure D- 88.  | Plot of pollution indicative species biomass at station SBC1 for 1999 through 2004. .    | D-146 |
| Figure D- 89.  | Plot of the benthic IBI at station SBD1 from 1999 through 2004. ....                     | D-147 |
| Figure D- 90.  | Plot of total benthic community abundance at station SBD1 for 1999 through 2004. . .     | D-148 |
| Figure D- 91.  | Plot of total benthic community biomass at station SBD1 for 1999 through 2004. . . .     | D-149 |
| Figure D- 92.  | Plot of the Shannon-Weiner diversity index at station SBD1 for 1999 through 2004. .      | D-150 |
| Figure D- 93.  | Plot of pollution sensitive species abundance at station SBD1 for 1999 through 2004. .   | D-151 |
| Figure D- 94.  | Plot of pollution indicative species abundance at station SBD1 for 1999 through 2004. D- | D-152 |
| Figure D- 95.  | Plot of pollution sensitive species biomass at station SBD1 for 1999 through 2004. . .   | D-153 |
| Figure D- 96.  | Plot of pollution indicative species biomass at station SBD1 for 1999 through 2004. .    | D-154 |
| Figure D- 97.  | Plot of the benthic IBI at station SBD2 from 1999 through 2004. ....                     | D-155 |
| Figure D- 98.  | Plot of total benthic community abundance at station SBD2 for 1999 through 2004. . .     | D-156 |
| Figure D- 99.  | Plot of total benthic community biomass at station SBD2 for 1999 through 2004. . . .     | D-157 |
| Figure D- 100. | Plot of the Shannon-Weiner diversity index at station SBD2 for 1999 through 2004. .      | D-158 |
| Figure D- 101. | Plot of pollution sensitive species abundance at station SBD2for 1999 through 2004. .    | D-159 |
| Figure D- 102. | Plot of pollution indicative species abundance at station SBD2 for 1999 through 2004. D- | D-160 |
| Figure D- 103. | Plot of pollution sensitive species biomass at station SBD2 for 1999 through 2004. . .   | D-161 |
| Figure D- 104. | Plot of pollution indicative species biomass at station SBD2 for 1999 through 2004. .    | D-162 |
| Figure D- 105. | Plot of the benthic IBI at station SBD4 from 1999 through 2004. ....                     | D-163 |
| Figure D- 106. | Plot of total benthic community abundance at station SBD4 for 1999 through 2004. . .     | D-164 |
| Figure D- 107. | Plot of total benthic community biomass at station SBD4 for 1999 through 2004. . . .     | D-165 |
| Figure D- 108. | Plot of the Shannon-Weiner diversity index at station SBD4 for 1999 through 2004. .      | D-166 |

|                |  |       |
|----------------|--|-------|
| Figure D- 109. | Plot of pollution sensitive species abundance at station SBD4 for 1999 through 2004. . | D-167 |
| Figure D- 110. | Plot of pollution indicative species abundance at station SBD4 for 1999 through 2004.  | D-168 |
| Figure D- 111. | Plot of pollution sensitive species biomass at station SBD4 for 1999 through 2004. . . | D-169 |
| Figure D- 112. | Plot of pollution indicative species biomass at station SBD4 for 1999 through 2004. .  | D-170 |






Figure D-1. Plot of the benthic IBI at station ELC1 from 1999 through 2004.

## ELC1




Figure D- 2. Plot of total benthic community abundance at station ELC1 for 1999 through 2004.

## ELC1




Figure D- 3. Plot of total benthic community biomass at station ELC1 for 1999 through 2004.




Figure D- 4. Plot of the Shannon-Weiner diversity index at station ELC1 for 1999 through 2004.




Figure D- 5. Plot of pollution sensitive species abundance at station ELC1 for 1999 through 2004.




Figure D- 6. Plot of pollution indicative species abundance at station ELC1 for 1999 through 2004.




Figure D- 7. Plot of pollution sensitive species biomass at station ELC1 for 1999 through 2004.

## ELC1




Figure D- 8. Plot of pollution indicative species biomass at station ELC1 for 1999 through 2004.






Figure D- 9. Plot of the benthic IBI at station ELD1 from 1999 through 2004.

## ELD1




Figure D- 10. Plot of total benthic community abundance at station ELD1 for 1999 through 2004.

## ELD1




Figure D- 11. Plot of total benthic community biomass at station ELD1 for 1999 through 2004.




Figure D- 12. Plot of the Shannon-Weiner diversity index at station ELD1 for 1999 through 2004.




Figure D- 13. Plot of pollution sensitive species abundance at station ELD1 for 1999 through 2004.

## ELD1




Figure D- 14. Plot of pollution indicative species abundance at station ELD1 for 1999 through 2004.

## ELD1




Figure D- 15. Plot of pollution sensitive species biomass at station ELD1 for 1999 through 2004.

## ELD1




Figure D- 16. Plot of pollution indicative species biomass at station ELD1 for 1999 through 2004.



## ELF1




Figure D- 17. Plot of the benthic IBI at station ELF1 from 1999 through 2004.

## ELF1




Figure D- 18. Plot of total benthic community abundance at station ELF1 for 1999 through 2004.

## ELF1




Figure D- 19. Plot of total benthic community biomass at station ELF1 for 1999 through 2004.




Figure D- 20. Plot of the Shannon-Weiner diversity index at station ELF1 for 1999 through 2004.




Figure D- 21. Plot of pollution sensitive species abundance at station ELF1 for 1999 through 2004.




Figure D- 22. Plot of pollution indicative species abundance at station ELF1 for 1999 through 2004.

## ELF1




Figure D- 23. Plot of pollution sensitive species biomass at station ELF1 for 1999 through 2004.




Figure D- 24. Plot of pollution indicative species biomass at station ELF1 for 1999 through 2004.






Figure D- 25. Plot of the benthic IBI at station EBB1 from 1999 through 2004.




Figure D- 26. Plot of total benthic community abundance at station EBB1 for 1999 through 2004.




Figure D- 27. Plot of total benthic community biomass at station EBB1 for 1999 through 2004.

## EBB1




Figure D- 28. Plot of the Shannon-Weiner diversity index at station EBB1 for 1999 through 2004.




Figure D- 29. Plot of pollution sensitive species abundance at station EBB1 for 1999 through 2004.




Figure D- 30. Plot of pollution indicative species abundance at station EBB1 for 1999 through 2004.

## EBB1




Figure D- 31. Plot of pollution sensitive species biomass at station EBB1 for 1999 through 2004.




Figure D- 32. Plot of pollution indicative species biomass at station EBB1 for 1999 through 2004.



## LFA1




Figure D- 33. Plot of the benthic IBI at station LFA1 from 1999 through 2004.




Figure D- 34. Plot of total benthic community abundance at station LFA1 for 1999 through 2004.

## LFA1




Figure D- 35. Plot of total benthic community biomass at station LFA1 for 1999 through 2004.




Figure D- 36. Plot of the Shannon-Weiner diversity index at station LFA1 for 1999 through 2004.




Figure D- 37. Plot of pollution sensitive species abundance at station LFA1 for 1999 through 2004.




Figure D- 38. Plot of pollution indicative species abundance at station LFA1 for 1999 through 2004.




Figure D- 39. Plot of pollution sensitive species biomass at station LFA1 for 1999 through 2004.




Figure D- 40. Plot of pollution indicative species biomass at station LFA1 for 1999 through 2004.



## LFB1




Figure D- 41. Plot of the benthic IBI at station LFB1 from 1999 through 2004.

## LFB1




Figure D-42. Plot of total benthic community abundance at station LFB1 for 1999 through 2004.

## LFB1




Figure D- 43. Plot of total benthic community biomass at station LFB1 for 1999 through 2004.

## LFB1




Figure D- 44. Plot of the Shannon-Weiner diversity index at station LFB1 for 1999 through 2004.

## LFB1




Figure D- 45. Plot of pollution sensitive species abundance at station LFB1 for 1999 through 2004.




Figure D- 46. Plot of pollution indicative species abundance at station LFB1 for 1999 through 2004.

## LFB1




Figure D- 47. Plot of pollution sensitive species biomass at station LFB1 for 1999 through 2004.




Figure D- 48. Plot of pollution indicative species biomass at station LFB1 for 1999 through 2004.






Figure D- 49. Plot of the benthic IBI at station WBB1 from 1999 through 2004.

## WBB1




Figure D- 50. Plot of total benthic community abundance at station WBB1 for 1999 through 2004.

## WBB1




Figure D- 51. Plot of total benthic community biomass at station WBB1 for 1999 through 2004.

## WBB1




Figure D- 52. Plot of the Shannon-Weiner diversity index at station WBB1 for 1999 through 2004.




Figure D- 53. Plot of pollution sensitive species abundance at station WBB1 for 1999 through 2004.




Figure D- 54. Plot of pollution indicative species abundance at station WBB1 for 1999 through 2004.

## WBB1




Figure D- 55. Plot of pollution sensitive species biomass at station WBB1 for 1999 through 2004.

## WBB1




Figure D- 56. Plot of pollution indicative species biomass at station WBB1 for 1999 through 2004.






Figure D- 57. Plot of the benthic IBI at station WBB5 from 1999 through 2004.




Figure D- 58. Plot of total benthic community abundance at station WBB5 for 1999 through 2004.

## WBB5




Figure D- 59. Plot of total benthic community biomass at station WBB5 for 1999 through 2004.




Figure D- 60. Plot of the Shannon-Weiner diversity index at station WBB5 for 1999 through 2004.




Figure D- 61. Plot of pollution sensitive species abundance at station WBB5 for 1999 through 2004.




Figure D- 62. Plot of pollution indicative species abundance at station WBB5 for 1999 through 2004.




Figure D- 63. Plot of pollution sensitive species biomass at station WBB5 for 1999 through 2004.

## WBB5




Figure D- 64. Plot of pollution indicative species biomass at station WBB5 for 1999 through 2004.






Figure D- 65. Plot of the benthic IBI at station SBA1 from 1999 through 2004.

## SBA1




Figure D- 66. Plot of total benthic community abundance at station SBA1 for 1999 through 2004.




Figure D- 67. Plot of total benthic community biomass at station SBA1 for 1999 through 2004.




Figure D- 68. Plot of the Shannon-Weiner diversity index at station SBA1 for 1999 through 2004.




Figure D- 69. Plot of pollution sensitive species abundance at station SBA1 for 1999 through 2004.




Figure D- 70. Plot of pollution indicative species abundance at station SBA1 for 1999 through 2004.




Figure D- 71. Plot of pollution sensitive species biomass at station SBA1 for 1999 through 2004.




Figure D- 72. Plot of pollution indicative species biomass at station SBA1 for 1999 through 2004.






Figure D- 73. Plot of the benthic IBI at station SBB1 from 1999 through 2004.




Figure D- 74. Plot of total benthic community abundance at station SBB1 for 1999 through 2004.




Figure D- 75. Plot of total benthic community biomass at station SBB1 for 1999 through 2004.




Figure D- 76. Plot of the Shannon-Weiner diversity index at station SBB1 for 1999 through 2004.




Figure D- 77. Plot of pollution sensitive species abundance at station SBB1 for 1999 through 2004.




Figure D- 78. Plot of pollution indicative species abundance at station SBB1 for 1999 through 2004.

## SBB1




Figure D- 79. Plot of pollution sensitive species biomass at station SBB1 for 1999 through 2004.




Figure D- 80. Plot of pollution indicative species biomass at station SBB1 for 1999 through 2004.






Figure D- 81. Plot of the benthic IBI at station SBC1 from 1999 through 2004.




Figure D- 82. Plot of total benthic community abundance at station SBC1 for 1999 through 2004.




Figure D- 83. Plot of total benthic community biomass at station SBC1 for 1999 through 2004.

## SBC1




Figure D- 84. Plot of the Shannon-Weiner diversity index at station SBC1 for 1999 through 2004.




Figure D- 85. Plot of pollution sensitive species abundance at station SBC1 for 1999 through 2004.




Figure D- 86. Plot of pollution indicative species abundance at station SBC1 for 1999 through 2004.




Figure D- 87. Plot of pollution sensitive species biomass at station SBC1 for 1999 through 2004.




Figure D- 88. Plot of pollution indicative species biomass at station SBC1 for 1999 through 2004.






Figure D- 89. Plot of the benthic IBI at station SBD1 from 1999 through 2004.




Figure D- 90. Plot of total benthic community abundance at station SBD1 for 1999 through 2004.




Figure D- 91. Plot of total benthic community biomass at station SBD1 for 1999 through 2004.




Figure D- 92. Plot of the Shannon-Weiner diversity index at station SBD1 for 1999 through 2004.




Figure D- 93. Plot of pollution sensitive species abundance at station SBD1 for 1999 through 2004.




Figure D- 94. Plot of pollution indicative species abundance at station SBD1 for 1999 through 2004.




Figure D- 95. Plot of pollution sensitive species biomass at station SBD1 for 1999 through 2004.




Figure D- 96. Plot of pollution indicative species biomass at station SBD1 for 1999 through 2004.






Figure D- 97. Plot of the benthic IBI at station SBD2 from 1999 through 2004.



Figure D- 98. Plot of total benthic community abundance at station SBD2 for 1999 through 2004.

## SBD2




Figure D- 99. Plot of total benthic community biomass at station SBD2 for 1999 through 2004.




Figure D- 100. Plot of the Shannon-Weiner diversity index at station SBD2 for 1999 through 2004.




Figure D- 101. Plot of pollution sensitive species abundance at station SBD2 for 1999 through 2004.




Figure D- 102. Plot of pollution indicative species abundance at station SBD2 for 1999 through 2004.




Figure D- 103. Plot of pollution sensitive species biomass at station SBD2 for 1999 through 2004.

## SBD2




Figure D- 104. Plot of pollution indicative species biomass at station SBD2 for 1999 through 2004.






Figure D- 105. Plot of the benthic IBI at station SBD4 from 1999 through 2004.

## SBD4




Figure D- 106. Plot of total benthic community abundance at station SBD4 for 1999 through 2004.

## SBD4




Figure D- 107. Plot of total benthic community biomass at station SBD4 for 1999 through 2004.




Figure D- 108. Plot of the Shannon-Weiner diversity index at station SBD4 for 1999 through 2004.




Figure D- 109. Plot of pollution sensitive species abundance at station SBD4 for 1999 through 2004.




Figure D- 110. Plot of pollution indicative species abundance at station SBD4 for 1999 through 2004.

## SBD4




Figure D- 111. Plot of pollution sensitive species biomass at station SBD4 for 1999 through 2004.




Figure D- 112. Plot of pollution indicative species biomass at station SBD4 for 1999 through 2004.